

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

GENFARM FLUTRIAFOL 500 SC FUNGICIDE

ACTIVE CONSTITUENT: 500 g/L FLUTRIAFOL

GROUP	3	FUNGICIDE
--------------	----------	------------------

**For control of certain fungal diseases on wheat, barley and
canola when mixed with fertiliser as per the Directions for
Use table**

Contents: 10L – 1000L

**Landmark Operations Limited
Suite 3, Building “B, 11 Talavera Road,
Macquarie Park NSW, 2113
Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92**

DIRECTIONS FOR USE: ALL STATES

Crop	Disease	Rate per hectare	WHP	Critical Comments																								
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	200mL	4 weeks Grazing	<p>Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties.</p> <p>Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 200mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide.</p> <table border="0"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr> <td>60</td> <td>333*</td> </tr> <tr> <td>70</td> <td>285</td> </tr> <tr> <td>80</td> <td>250</td> </tr> <tr> <td>100</td> <td>200</td> </tr> <tr> <td>120</td> <td>166</td> </tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)	60	333*	70	285	80	250	100	200	120	166												
Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)																											
60	333*																											
70	285																											
80	250																											
100	200																											
120	166																											
Wheat	Stripe rust (<i>Puccinia striiformis</i>)	100 or 200mL		<p>For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected.</p> <p>Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide.</p> <table border="0"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th colspan="2">Volume of product per 100kg of fertiliser (mL)</th> </tr> <tr> <th></th> <th>100mL rate</th> <th>200mL rate</th> </tr> </thead> <tbody> <tr> <td>50</td> <td>200</td> <td>400 *</td> </tr> <tr> <td>60</td> <td>166</td> <td>333 *</td> </tr> <tr> <td>70</td> <td>143</td> <td>285</td> </tr> <tr> <td>80</td> <td>125 †</td> <td>250</td> </tr> <tr> <td>100</td> <td>100 †</td> <td>200</td> </tr> <tr> <td>120</td> <td>88 †</td> <td>166</td> </tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)			100mL rate	200mL rate	50	200	400 *	60	166	333 *	70	143	285	80	125 †	250	100	100 †	200	120	88 †	166
	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)																										
	100mL rate	200mL rate																										
50	200	400 *																										
60	166	333 *																										
70	143	285																										
80	125 †	250																										
100	100 †	200																										
120	88 †	166																										
	Septoria tritici blotch (<i>S. tritici</i>) Take-all (<i>Geaumannomyces graminis</i>)	200mL																										
Barley	Powdery mildew (<i>Erysiphe graminis</i>) Scald (<i>Rhynchosporium secalis</i>)	100 or 200mL	<p>* Application volumes over 300mL per 100kg of fertiliser may adversely affect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers.</p> <p>† Where application volume is low, product may be diluted with water to improve coverage.</p>																									

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

APPLIED IN-FURROW AS A FERTILISER TREATMENT

CANOLA, WHEAT AND BARLEY - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING

GENERAL INSTRUCTIONS

Fertiliser Application

Ensure that treated fertiliser is dry before sowing with the crop.

Sowing: Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered.

Do not mix seed with treated fertiliser.

FUNGICIDE RESISTANCE WARNING

GROUP	3	FUNGICIDE
-------	---	-----------

Genfarm Flutriafol 500 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Genfarm Flutriafol 500 SC Fungicide is a Group 3 fungicide.

Some naturally occurring individual fungi resistant to Genfarm Flutriafol 500 SC Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly.

These resistant fungi will not be controlled by Genfarm Flutriafol 500 SC Fungicide or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Landmark Operations Limited accepts no liability for any losses that may result from the failure of Genfarm Flutriafol 500 SC Fungicide to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the original container, tightly closed in a safe well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight.

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple rinse container for disposal. Dispose of rinsate by adding it to the spray tank. DO NOT dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection point or similar container management program site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers and product.

For REFILLABLE containers (110L, 1000L): Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Harmful if absorbed by skin contact. May irritate the skin. Will irritate the eyes. Avoid contact with eyes and skin. When opening the container and preparing product for use and using the product wear cotton overalls buttoned to the neck and wrist [or equivalent clothing], elbow-length chemical resistant gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each days use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Landmark Operations Limited. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Landmark Operations Limited accepts no responsibility for any consequences resulting from

the use of this product. Landmark Operations Limited will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

Batch No:

Date of Manufacture:

APVMA Approval No: 67202/100097

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS AUSTRALIA WIDE
--	---

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Harmful if inhaled; Avoid breathing spray; Do not eat, drink or smoke when using this product; If inhaled remove person to fresh air and keep comfortable for breathing