

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Genfarm
TRIADIMEFON 500 WP
Fungicide

ACTIVE CONSTITUENT:
500 g/kg TRIADIMEFON

GROUP	C	FUNGICIDE
--------------	----------	------------------

Wheat:

For control of leaf rust, septoria and speckled leaf blotch.

Barley:

For control of powdery mildew

MEASURE PACK CONTAINS:

1 Kg NET

NOT TO BE SOLD SEPARATELY

BEFORE USE READ ALL DIRECTIONS ON THE OUTER PACK

DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS

APVMA Approval No: 58551/1/0304

Genfarm Crop Protection Pty Ltd
Suite 3, Level 1, 64 Talavera Road,
Macquarie Park,
NSW, 2113
Tel: (02) 9889 5400

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Genfarm
TRIADIMEFON 500 WP
Fungicide

ACTIVE CONSTITUENT:
500 g/kg TRIADIMEFON

GROUP	C	FUNGICIDE
--------------	----------	------------------

Wheat:

For control of leaf rust, septoria and speckled leaf blotch.

Barley:

For control of powdery mildew

MEASURE PACK CONTAINS:

(5 kg Net)

NOT TO BE SOLD SEPARATELY

BEFORE USE READ ALL DIRECTIONS ON THE OUTER PACK

DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS

APVMA Approval No: 58551/5/0105

Genfarm Crop Protection Pty Ltd
Suite 3, Level 1, 64 Talavera Road,
Macquarie Park,
NSW, 2113
Tel: (02) 9889 5400

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Genfarm

TRIADIMEFON 500 WP
Fungicide

ACTIVE CONSTITUENT:
500 g/kg TRIADIMEFON

GROUP	C	FUNGICIDE
--------------	----------	------------------

Wheat:

For control of leaf rust, septoria and speckled leaf blotch.

Barley:

For control of powdery mildew

10 Kg NET

PRIMARY PACK CONTAINS: 2 x 5kg MEASURE PACKS
WHICH IT IS ILLEGAL TO SELL SEPARATELY

DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS

Genfarm Crop Protection Pty Ltd

Suite 3, Level 1, 64 Talavera Road,

Macquarie Park,

NSW, 2113

Tel: (02) 9889 5400

DIRECTIONS FOR USE

Crop	Disease/ Pest	State	Rate	Critical Comments
Barley	Powdery mildew.	NSW, VIC, SA, WA ONLY	250 g/ha	Apply at the first signs of stress
Wheat	Leaf Rust (Puccinia recondita)	NSW, VIC, SA, WA ONLY	125 - 250 g/ha	For spray timings see general instructions. Normally use 125 g/ha. Only use 250g/ha where the yield is expected to be greater than 4-5 tonnes/ha or if a longer residual is required.
	Speckled leaf blotch.	Mid and North Western Victoria	250 g/ha	Where this disease commonly affects the Flag leaf, apply to susceptible varieties at the 5-6 tiller stage. Re-infestation can occur if moist conditions prevail
	Septoria	Sthn NSW, Vic only	125 - 250 g/ha	For control of light to moderate infections. Applications should coincide with early head emergence. Tank mixes with Benomyl may improve control in severe infections.

Applied as a fertilizer treatment

Crop	Disease/ Pest	State	Rate	Critical Comments
Wheat	Leaf Rust (Puccinia recondita)	NSW, VIC, SA, WA ONLY	200 g/ha	Use the higher rate where conditions are conducive to very high rates of disease pressure. Also in years when the incidence of disease is expected to be high as forecast by Department of Agriculture or reputable disease forecasters. Application: Apply product evenly onto the fertilizer. If using DAP or similar granulated product, use as powder. If adding to a particularly dusty – dry fertilizer granule, dilute with minimum water (up to a maximum of 8 Litres) ensure constant mixing of the suspension, using a “granny pot” or similar device. Calculation of the rate required / tonne fertiliser is below $1000 \div (\text{fert rate} / \text{ha}) \times \text{rate of fungicide (kg)}$ e.g. using 100 kg fert / ha, and wanting 200 gm/ha fungicide; $1000 \div 100 \times 0.2 = 2 \text{ kg / tonne.}$

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

DO NOT HARVEST FOR 3 MONTHS AFTER APPLICATION

GENERAL INSTRUCTIONS

Genfarm Triadimefon 500 WP Fungicide may be used by Aircraft or Ground spraying equipment.

Spray timings for rust control: Determine the susceptibility of the Wheat or Barley variety, e.g. Department Agriculture data, to determine spray requirements.

Seedling infections: When approx 20% of leaves show signs of infection during tillering or jointing, spray within a week

Adult infections:

Susceptible varieties: Spray within one week if approx 10% of leaves show infection.

Do not delay.

Moderately Susceptible varieties: Spray within one week if approx 15 - 20% of leaves show infection. Do not delay.

Moderately Resistant and Resistant varieties: Monitor vigilantly. Spray within one week if rust appears and spreads.

FUNGICIDE RESISTANCE WARNING:

GROUP	C	FUNGICIDE
-------	----------	-----------

Genfarm Triadimefon 500 WP Fungicide is a member of the D.M.I group of fungicides. For fungicide resistance management the product is a Group C fungicide.

Some naturally occurring individual fungi resistant to the product and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Genfarm Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

MIXING

The required quantity of Genfarm Triadimefon 500 WP Fungicide should be thoroughly mixed with a small quantity of clean, filtered water to make a creamy mix. Add this blend to the required quantity of water in the tank and mix thoroughly. Agitate prior to and during spraying.

DO NOT mix more than can be used for the day's spraying.

DO NOT tank mix with trace elements or other chemicals without first checking compatibility.

APPLICATION

Ground Rig: Use a convenient amount of water, e.g. 30 - 100 L water / ha

Aircraft: Apply at not less than 10 L / ha, preferably in a cross wind of greater than 5 knots. If this is not possible, increase the volume to 30 L / ha and increase the droplet size. Spray in the morning and reduce the swathe width. Booms, with fan nozzles set forward at 45° forward to the airstream and Micronairs with blade settings of 30 - 50° are equally satisfactory. Use proper measures for gauging wind speed, direction and variability of the wind.

PRECAUTIONS

Feed treated with this product must not be used for animal consumption, poultry feed or mixed with animal feed.

DO NOT mix leaves treated with this product with feed intended for animal consumption.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed treated leaf matter to wild or domestic birds or other animals.

Harmful to fish.

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store the unused product in the closed original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Do not allow seed treated with this product to contaminate grain intended for human consumption.

Do not use seed treated with this product for human consumption.

Bags which have held treated seed with this product are not to be used for any other purpose.

Shake empty bag. If using as a liquid, single rinse the bag before disposal and add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. Puncture and bury empty containers in a local authority landfill. If no landfill available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS:

Poisonous if swallowed. May irritate the eyes and skin. Do not inhale dust. When preparing product for use wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and disposable dust mask. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. (Phone Australia: 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information see Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

For specialist advice in an emergency dial

1 800 033 111

24 hours Australia wide.

Genfarm Crop Protection Pty Ltd

Suite 3, Level 1, 64 Talavera Road,

Macquarie Park,

NSW, 2113

Tel: (02) 9889 5400

Batch No.:

Date of Manufacture:

APVMA Approval No: 58551/10A/0105

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Genfarm

TRIADIMEFON 500 WP
Fungicide

ACTIVE CONSTITUENT:
500 g/kg TRIADIMEFON

GROUP	C	FUNGICIDE
--------------	----------	------------------

Wheat:

For control of leaf rust, septoria and speckled leaf blotch.

Barley:

For control of powdery mildew

10 Kg NET

PRIMARY PACK CONTAINS: 10 x 1kg MEASURE PACKS
WHICH IT IS ILLEGAL TO SELL SEPARATELY

DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS

Genfarm Crop Protection Pty Ltd

Suite 3, Level 1, 64 Talavera Road,

Macquarie Park,

NSW, 2113

Tel: (02) 9889 5400

DIRECTIONS FOR USE

Crop	Disease/ Pest	State	Rate	Critical Comments
Barley	Powdery mildew.	NSW, VIC, SA, WA ONLY	250 g/ha	Apply at the first signs of stress
Wheat	Leaf Rust (Puccinia recondita)	NSW, VIC, SA, WA ONLY	125 - 250 g/ha	For spray timings see general instructions. Normally use 125 g/ha. Only use 250g/ha where the yield is expected to be greater than 4-5 tonnes/ha or if a longer residual is required.
	Speckled leaf blotch.	Mid and North Western Victoria	250 g/ha	Where this disease commonly affects the Flag leaf, apply to susceptible varieties at the 5-6 tiller stage. Re-infestation can occur if moist conditions prevail
	Septoria	Sthn NSW, Vic only	125 - 250 g/ha	For control of light to moderate infections. Applications should coincide with early head emergence. Tank mixes with Benomyl may improve control in severe infections.

Applied as a fertilizer treatment

Crop	Disease/ Pest	State	Rate	Critical Comments
Wheat	Leaf Rust (Puccinia recondita)	NSW, VIC, SA, WA ONLY	200 g/ha	Use the higher rate where conditions are conducive to very high rates of disease pressure. Also in years when the incidence of disease is expected to be high as forecast by Department of Agriculture or reputable disease forecasters. Application: Apply product evenly onto the fertilizer. If using DAP or similar granulated product, use as powder. If adding to a particularly dusty – dry fertilizer granule, dilute with minimum water (up to a maximum of 8 Litres) ensure constant mixing of the suspension, using a “granny pot” or similar device. Calculation of the rate required / tonne fertiliser is below $1000 \div (\text{fert rate} / \text{ha}) \times \text{rate of fungicide (kg)}$ e.g. using 100 kg fert / ha, and wanting 200 gm/ha fungicide; $1000 \div 100 \times 0.2 = 2 \text{ kg / tonne.}$

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

DO NOT HARVEST FOR 3 MONTHS AFTER APPLICATION

GENERAL INSTRUCTIONS

Genfarm Triadimefon 500 WP Fungicide may be used by Aircraft or Ground spraying equipment.

Spray timings for rust control: Determine the susceptibility of the Wheat or Barley variety, e.g. Department Agriculture data, to determine spray requirements.

Seedling infections: When approx 20% of leaves show signs of infection during tillering or jointing, spray within a week

Adult infections:

Susceptible varieties: Spray within one week if approx 10% of leaves show infection.

Do not delay.

Moderately Susceptible varieties: Spray within one week if approx 15 - 20% of leaves show infection. Do not delay.

Moderately Resistant and Resistant varieties: Monitor vigilantly. Spray within one week if rust appears and spreads.

FUNGICIDE RESISTANCE WARNING:

GROUP	C	FUNGICIDE
-------	----------	-----------

Genfarm Triadimefon 500 WP Fungicide is a member of the D.M.I group of fungicides. For fungicide resistance management the product is a Group C fungicide.

Some naturally occurring individual fungi resistant to the product and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Genfarm Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

MIXING

The required quantity of Genfarm Triadimefon 500 WP Fungicide should be thoroughly mixed with a small quantity of clean, filtered water to make a creamy mix. Add this blend to the required quantity of water in the tank and mix thoroughly. Agitate prior to and during spraying.

DO NOT mix more than can be used for the day's spraying.

DO NOT tank mix with trace elements or other chemicals without first checking compatibility.

APPLICATION

Ground Rig: Use a convenient amount of water, e.g. 30 - 100 L water / ha

Aircraft: Apply at not less than 10 L / ha, preferably in a cross wind of greater than 5 knots. If this is not possible, increase the volume to 30 L / ha and increase the droplet size. Spray in the morning and reduce the swathe width. Booms, with fan nozzles set forward at 45° forward to the airstream and Micronairs with blade settings of 30 - 50° are equally satisfactory. Use proper measures for gauging wind speed, direction and variability of the wind.

PRECAUTIONS

Feed treated with this product must not be used for animal consumption, poultry feed or mixed with animal feed.

DO NOT mix leaves treated with this product with feed intended for animal consumption.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed treated leaf matter to wild or domestic birds or other animals.

Harmful to fish.

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store the unused product in the closed original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Do not allow seed treated with this product to contaminate grain intended for human consumption.

Do not use seed treated with this product for human consumption.

Bags which have held treated seed with this product are not to be used for any other purpose.

Shake empty bag. If using as a liquid, single rinse the bag before disposal and add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. Puncture and bury empty containers in a local authority landfill. If no landfill available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS:

Poisonous if swallowed. May irritate the eyes and skin. Do not inhale dust. When preparing the product for use wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and disposable dust mask. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. (Phone Australia: 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information see Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

For specialist advice in an emergency dial

1 800 033 111

24 hours Australia wide.

Genfarm Crop Protection Pty Ltd

Suite 3, Level 1, 64 Talavera Road,

Macquarie Park,

NSW, 2113

Tel: (02) 9889 5400

Batch No.:

Date of Manufacture:

APVMA Approval No: 58551/10/0105