

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

GENFARM TRIADIMENOL T FLOWABLE SEED DRESSING

**ACTIVE CONSTITUENTS: 150 g/L TRIADIMENOL
4 g/L TRIFLUMURON**

GROUP	3	FUNGICIDE
--------------	----------	------------------

GROUP	15A	INSECTICIDE
--------------	------------	--------------------

For the control or suppression of various diseases of wheat, barley and oats as specified in the Directions For Use Table

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

Contents: 5L, 10L, 20L, 100, 110, 200L

Landmark Operations Limited
Suite 2, Level 3, 64 Talavera Road,
North Ryde,
NSW, 2113
Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92

STORAGE AND DISPOSAL

Product and original container

Store in closed original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Treated Seed and Containers of Treated Seed

When treated seed is stored it should be kept apart from other grain and the bags or other containers should be clearly marked to indicate the contents have been treated with this product.

DO NOT use treated seed for human consumption. Bags which have held treated seed are not to be used for any other purpose.

Application Equipment

Rinse all equipment with clean water immediately after use and dispose of rinsings in a dedicated disposal pit, specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

CONDITIONS OF SALE

The use of this product is beyond the control of Landmark Operations Limited. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Landmark Operations Limited accepts no responsibility for any consequences resulting from the use of this product. Landmark Operations Limited will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

APVMA Approval No: 66315/53368

Batch No:

Date of Manufacture:

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS AUSTRALIA WIDE
--	---



CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

GENFARM TRIADIMENOL T FLOWABLE SEED DRESSING

**ACTIVE CONSTITUENTS: 150 g/L TRIADIMENOL
4 g/L TRIFLUMURON**

GROUP	3	FUNGICIDE
--------------	----------	------------------

GROUP	15A	INSECTICIDE
--------------	------------	--------------------

For the control or suppression of various diseases of wheat, barley and oats as specified in the Directions
For Use Table

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

APVMA Approval No: 66315/53368

Landmark Operations Limited
Suite 2, Level 3, 64 Talavera Road,
North Ryde,
NSW, 2113
Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92

DIRECTIONS FOR USE

Restraint:

DO NOT treat physically damaged seed.

Crop	Pest/Disease	Rate/ha	Critical Comments
Wheat	Bunt, flag smut (seed and soil borne), loose smut, suppression of seedling infections of stripe rust.	100 mL/100 kg of seed	<p>Genfarm Triadimenol T Flowable Seed Dressing should be mixed with water to give even coverage of seed. The quantity of water used for mixing will vary depending on type of equipment and quality of seed. Use a minimum of 400 mL of mixture (ie Triadimenol T plus water) with each 100 kg of seed. Do not use more than 600 mL of mixture product + water) with each 100 kg of seed. Whatever dilution is used it is essential that 100 mL (Or 150 mL) of Triadimenol T is applied per 100 kg of seed. The mixture should be gently stirred regularly. If nasal irritation occurs use a face mask while treating seed.</p> <p>Use 150mL/1000kg seed rate where foliar diseases occur frequently and where any of the following are required:</p> <ul style="list-style-type: none"> - Longer suppression of stripe rust on seedlings - Control of seedling infections of powdery mildew - Suppression of speckled leaf blotch in wheat - Extended suppression of leaf scald in barley: and use higher rate on susceptible varieties or where disease pressure is heavy.
Oats	Covered and loose smut		
Barley	Covered and loose smut. Suppression of seedling infections of powdery mildew.		
	Suppression of seedling infections of leaf scald (not Qld)		
Wheat	Bunt, flag smut (seed and soil borne), loose smut, suppression of seedling infections of stripe rust.	150 mL/ 100 kg seed	<ul style="list-style-type: none"> - Suppression of speckled leaf blotch in wheat - Extended suppression of leaf scald in barley: and use higher rate on susceptible varieties or where disease pressure is heavy.
	Suppression of seedling infections of speckled leaf blotch (NOT Qld and WA)		
Barley	Covered and loose smut. Suppression of seedling infections of powdery mildew.		
	Suppression of seedling leaf scald (NOT Qld)		
Wheat, barley, oats	Protection against insect pests of stored grains: granary weevil, lesser grain borer, rice weevil, rust-red flour beetle, saw-toothed grain beetle	100 or 150 mL/kg seed	Both rates protect treated seed from grain insect pests. Use rate appropriate to disease control required.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING: DO NOT GRAZE PLANTS GROWN FROM TREATED SEED OR CUT FOR STOCK FOOD WITHIN 5 WEEKS OF SOWING

GENERAL INSTRUCTIONS

Wheat

Genfarm Triadimenol T Flowable Seed Dressing, especially at the higher rate of 150 mL/100kg seed, may delay emergence of wheat. Crops will recover rapidly in most soil types. However, seed treated with 150mL/100kg seed rate should not be sown in soils that are likely to crust or clod at the surface after sowing. Sowing Genfarm Triadimenol T treated seed deeper than 50 mm may also slightly reduce percentage plant establishment in heavier soils and sowing rates may need to be adjusted accordingly. The possible delay in emergence with the 150 mL/100kg seed rate may counteract the advantage of suppression of speckled leaf blotch.

Fungicide Resistance Warning

GROUP

3

FUNGICIDE

Genfarm Triadimenol T Flowable Seed Dressing is a member of the DMI group of fungicides. For fungicide resistance management Genfarm Triadimenol T Flowable Seed Dressing is a Group 3 fungicide.

Some naturally occurring individual fungi resistant to Genfarm Triadimenol T Flowable Seed Dressing and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Genfarm Triadimenol T Flowable Seed Dressing or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Landmark Operations Limited accepts no liability for any losses that may result from the failure of Genfarm Triadimenol T Flowable Seed Dressing to control resistant fungi.

Insecticide Resistance Warning

GROUP

15A

INSECTICIDE

For insecticide resistance management Genfarm Triadimenol T Flowable Seed Dressing is a Group 15A insecticide. Some naturally occurring insect biotypes resistant to Genfarm Triadimenol T Flowable Seed Dressing and other Group 15A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Genfarm Triadimenol T Flowable Seed Dressing and other Group 15A insecticides are used repeatedly. The effectiveness of Genfarm Triadimenol T Flowable Seed Dressing on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Landmark Operations Limited accepts no liability for any losses that may result from the failure of Genfarm Triadimenol T Flowable Seed Dressing to control resistant insects. Genfarm Triadimenol T Flowable Seed Dressing may be subject to specific resistance management strategies. For further information contact your local supplier, Landmark representative or local agricultural department agronomist.

Seed Quality

Genfarm Triadimenol T Flowable Seed Dressing should not be used on seed with more than 12% moisture content, or in sprung, sprouted, damaged or severely pinched seed, or seed of poor viability. If in doubt, have a germination test carried out on the seed before treatment to ensure that is of an acceptable standard.

Treating seed with Genfarm Triadimenol T Flowable Seed Dressing as directed may raise the moisture level of the seed by up to 0.6%, depending on conditions at treatment. The use of Genfarm Triadimenol T Flowable Seed Dressing at the recommended rates will have no effect on the storage life of sound seed.

Storage of Treated Seed

Treated seed may be held over for sowing in the following season. If so, an additional insecticide treatment will be necessary. Treated seeds prevents insects from multiplying and preserves grain condition but may not kill adult insects migrating from other infected grain. If seed is not used immediately after treatment is should be stored in a dry, well ventilated place. Although Genfarm Triadimenol T Flowable Seed Dressing has no effect on the viability of treated seed, subsequent germination can be adversely affected by poor storage conditions such as high moisture combined with high temperatures. No liability can therefore be accepted for the performance of stored treated seed.

Cleaning Up

Equipment should be thoroughly cleaned with water. In the event of changing from this water based flowable formulation to a solvent-based liquid formulation, it is essential to rinse equipment with methylated spirits after thoroughly cleaning with water. If changing from a solvent based liquid to this water based flowable, thoroughly clean with methylated spirits then rinse with water. Failure to do so will result in clogging of equipment.

PRECAUTION

DO NOT allow seed treated with this product to contaminate seed intended for human consumption. If possible use a separate auger for treating seed. If using the same auger for treating seed and moving grain for human consumption, remove all residues of Genfarm Triadimenol T residues from the auger and avoid contaminating untreated seed with seed dressing residues.

PROTECTION OF LIVESTOCK

Seed treated with this product must not be used for animal consumption or poultry feed or mixed with animal feed. DO NOT allow seed treated with this product to contaminate seed intended for animal consumption.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed treated seed or otherwise expose to wild or domestic birds.

Harmful to fish and aquatic arthropods. DO NOT contaminate dams, rivers, ponds, waterways or drains with this product, used containers or bags which have held the treated seed.

STORAGE AND DISPOSAL

Product and original container

Store in closed original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Treated Seed and Containers of Treated Seed

When treated seed is stored it should be kept apart from other grain and the bags or other containers should be clearly marked to indicate the contents have been treated with this product.

DO NOT use treated seed for human consumption. Bags which have held treated seed are not to be used for any other purpose.

Application Equipment

Rinse all equipment with clean water immediately after use and dispose of rinsings in a dedicated disposal pit, specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

CONDITIONS OF SALE

The use of this product is beyond the control of Landmark Operations Limited. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Landmark Operations Limited accepts no responsibility for any consequences resulting from the use of this product. Landmark Operations Limited will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

Not classified as Hazardous according to the Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia criteria.
--