

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## GENFARM DIQUAT 200 NON-RESIDUAL HERBICIDE

ACTIVE CONSTITUENT: 200 g/L DIQUAT  
present as DIQUAT DIBROMIDE MONOHYDRATE

GROUP	L	HERBICIDE
-------	---	-----------

For pre-harvest crop desiccation and the control of a wide range of broadleaf weeds in certain crops as per Directions for Use.  
For application through aircraft and ground equipment.

**IMPORTANT: READ ATTACHED LEAFLET BEFORE USING THIS PRODUCT**

Contents: 20 L, 110 L, 1000 L

Nutrien Ag Solutions Limited  
Level 5, Building A, 26 Talavera Road,  
Macquarie Park, NSW 2113.  
Tel: (02) 9889 5400  
Product Support Tel: 1800 44 88 92

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or a place away from children, animals, food, feedstuffs, seed and fertilisers.

This containers can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple or preferably pressure rinse containers before disposal. Dispose of rinsate by adding it to the spray tank. DO NOT dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection point or similar container management program site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

*For Refillable containers (110L, 1000L):* Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## SAFETY DIRECTIONS

Very dangerous. Poisonous if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves, face shield or goggles, and half-face respirator or disposable respirator.

If clothing becomes contaminated with product or wet with spray, remove clothing immediately. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles, respirator and if rubber wash with detergent and warm water, and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 131126).

If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

## SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from the supplier.

## CONDITIONS OF SALE

The use of this product is beyond the control of Nutrien Ag Solutions Limited. Any provisions or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Nutrien Ag Solutions Limited does not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.


APVMA Approval No: 64889/58177

Batch Number:

Date of Manufacture:

Nutrien Ag Solutions Limited  
Level 5, Building A, 26 Talavera Road,  
Macquarie Park, NSW 2113.  
Tel: (02) 9889 5400  
Product Support Tel: 1800 44 88 92

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Causes damage to organs through prolonged or repeated exposure; If inhaled remove person to fresh air and keep comfortable for breathing; If skin or eye irritation persists get medical attention; Use only outdoors or in a well-ventilated area; Contaminated work clothing should not be allowed out of the workplace.
---

<b>BIPYRIDILIUM PESTICIDES, LIQUID, TOXIC (CONTAINS DIQUAT)</b>	
<b>UN NO. 3016</b>	<b>PG III</b>
<b>HAZCHEM 2X</b>	
<b>IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE</b>	
<b>FOR SPECIALIST ADVICE IN AN EMERGENCY 1800 033 111 24 HOURS AUSTRALIA WIDE</b>	

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## GENFARM DIQUAT 200 NON-RESIDUAL HERBICIDE

ACTIVE CONSTITUENT: 200 g/L DIQUAT  
present as DIQUAT DIBROMIDE MONOHYDRATE

GROUP	L	HERBICIDE
-------	---	-----------

For pre-harvest crop desiccation and the control of a wide range of broadleaf weeds in certain crops as per Directions for Use.  
For application through aircraft and ground equipment.

**IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT**

**APVMA Approval No.: 64889/58177**

**Nutrien Ag Solutions Limited  
Level 5, Building A, 26 Talavera Road,  
Macquarie Park, NSW 2113.  
Tel: (02) 9889 5400  
Product Support Tel: 1800 44 88 92**

**DIRECTIONS FOR USE****RESTRAINTS**

Best results may be obtained when application is made in dull weather or at the end of the day.

**DO NOT** spray when weeds are under drought stress or when covered with dust or soil.

**Pre-harvest crop desiccation:**

Crop	State	Rate *	WHP	Critical Comments	
Cotton (short stapled varieties only)	Qld, NSW, and WA only	2 or 3 L/ha	NIL	Apply when 85% of the bolls are open and remaining bolls are mature. Genfarm Diquat 200 can damage green bolls.	
Dry Beans, Peas, Lentils, Chickpeas, Faba beans	All States	2 or 3 L/ha	NIL	Spray as soon as the crop has reached full maturity. Helps overcome slow and uneven ripening and weed problems at harvest.	
Linseed	All States	2 or 3 L/ha	1 day grazing	Spray when the majority of seed heads are mature – 90 to 95% of seed heads have changed from yellow to brown and the seeds rattle inside the bolls. Desiccation reduces the period from maturity to harvest, particularly under wet or humid conditions.	
Lupins	All States	2 or 3 L/ha	1 day grazing	Spray as soon as crop has reached full maturity. Helps overcome slow and uneven ripening and weed problems at harvest.	
Mung Beans	NSW, Qld only	2 or 3 L/ha	nil	Apply when 80 to 90% of pods are black or brown. Desiccation of weeds and foliage aids timely and efficient harvesting, reduces harvester wear and tear but can increase harvest losses. Harvest 2 to 5 days after spraying.	
Perennial Legume Seed Crops	All States	1.5 to 3 L/ha	nil	<p><b>Lucerne</b> – Spray when 60 to 70% of the pods are brown/bluish and the seeds are yellow/brown and easily released from the pods.</p> <p><b>Red Clover</b> – Spray when majority of seed heads are brown and the seed is purple.</p> <p><b>White Clover</b> – Spray when majority of seeds are hard and yellow.</p>	The use of Genfarm Diquat 200 enables direct harvesting instead of cutting and windrowing and may result in higher seed quality. Harvest 3 to 4 days after spraying.
Pigeon Peas	Qld, NSW only	2 or 3 L/ha	4 days	Spray as soon as the crop has reached full maturity.	
Poppies	Tas only	3 to 4 L/ha	14 days	Spray after the poppies have reached the stripy capsule stage. Helps overcome slow and uneven ripening and weed problems at harvest.	
Potato (Haulm desiccation)	All States	3 to 4 L/ha	7 days	Apply as soon as crop is ready to harvest. <b>DO NOT</b> apply during drought periods, particularly when the tops will wilt during the day. In such conditions, wait at least 3 days after the soil has been well moistened by rain or irrigation. Leaf kill is rapid following spraying and usually complete within 4 days. Stem kill may take 10 to 14 days. Lift when desiccation is complete but where possible, wait for 14 days after spraying to allow skin to harden off. Use high water volumes to obtain coverage of dense haulm.	

Ground stored - pre-harvest weed control		1.5 L/ha plus 1.2 L Paraquat 250	7 days	To remove weed growth and facilitate digging, spray about 7 days prior to harvest. Where digging has been postponed and tubers stored in the ground often for a lengthy period, weed growth can be heavy and impede mechanical diggers unless removed.
--	--	--	--------	--

**Pre-harvest crop desiccation (continued):**

Crop	State	Rate *	WHP	Critical Comments
Canola	All States	1.5 or 3 L/ha	4 days	Spray when 70% of the pods are yellow and the seeds are brown/bluish and pliable. Canola ripens unevenly and is prone to pod shatter and seed loss. Direct harvest 4 to 7 days after spraying.
Rice	Qld, NSW only	2 to 3 L/ha	5 days	Spray when the grain is mature – not more than 2 to 3% of the grain is still at the milky stage and the grain moisture content must be less than 25%.
Sorghum	Qld, NSW only	2 to 3 L/ha	1 day grazing	Spray as soon as the seed is mature and the moisture content about 25%. Genfarm Diquat 200 will advance harvest and reduce seed losses due to differential ripening, seed shedding and birds.
Soybeans	Qld, NSW, WA only	2 to 3 L/ha	4 days	Spray when 80% of the pods are yellow/brown and the seeds are ripe – yellow and pliable. Desiccation of weeds and foliage aids timely and efficient harvesting, minimises cost and increases yields. Harvest 4 to 7 days after spraying.
Sugarcane	Qld, NSW only	2 to 3 L/ha	4 days	Spray all accessible faces a few days prior to burning to a depth of about 30 metres. The sprayed cane and weed growth quickly dries out and ensures a good burn and removal of trash prior to harvest.
		High volume hand spraying 200 mL/200L water		Spray to visible wetness.
Sunflower	All States	2 to 3 L/ha	4 days	Spray when the seed is mature, seed moisture 35% and below, kernel full and firm, the disc spongy when broken, florets loose and bracts browning off. Harvesting can commence as soon as vegetative parts of crop are desiccated, usually 7 to 14 days after spraying.
Sweet Potatoes	Qld only	3 to 4 L/ha	14 days	Apply 2 weeks prior to harvesting.

**General weed control**

Crop	Weeds Controlled	State	Rate *	WHP	Critical Comments
Aquatic areas	Duck Weeds, Red Azolla, Water Hyacinth, Salvinia, Marsilea, Water Lilies and Water Lettuce	All States	5 or 10 L/ha	10 days	Apply as an overall spray wetting foliage thoroughly. Clear water is necessary for best results as suspended soil particles interfere with herbicidal action. Use the higher rate for heavy infestations or for deep or dirty water
			400 mL plus 150 mL Agral per 100 L water		Small areas – spray to wet weeds thoroughly. About 1 mL of product should be sufficient to treat about 1 m <sup>2</sup> .
	Cattail and Pond Weeds		5 L/ megalitre water		Apply by injection below the surface or as a surface spray.
Asparagus	Broadleaf Weeds	All States	1.4 L/ha plus 800 mL Agral in 400 L water/ha	nil	Apply to control seedling weeds before spears have emerged.

Crop	Weeds Controlled	State	Rate *	WHP	Critical Comments	
Hops	Annual broadleaf and grass weeds	Vic and Tas only	700 mL to 1.4 L/ha may be mixed with 1.2 to 1.6 L/ha Paraquat 250 and/or 1.1 kg Simazine 900 Granules	nil	Apply as a directed inter-row spray prior to crop emerging from winter dormancy, using a minimum of 250 L/ha spray volume to ensure good and even coverage of weeds. <del>Add Wetter 1000 at the rate of 200mL/100 water — see General Instructions.</del> If used with Paraquat 250 Herbicide observe safety directions for that product.	
Infested areas	Cotton Thistle ( <i>Onopordum acanthium</i> )	Tas only	300 mL/ha plus 150 mL Agral in 100 L water	nil	Spot spray at the rosette stage before the centre shoot is 15cm tall. The spray should be applied to give complete wetting of the leaf surface. <b>DO NOT</b> use a lower rate or treat at a later growth stage.	
	Saffron Thistle	All States	2.8 L/ha plus 1 L Agral in 200 L water/ha		Apply as an overall treatment to prevent seeding.	
			100 mL plus 70 mL Agral per 15 L knapsack		Alternatively spot spray on the same basis.	
Lucerne	Capeweed and <i>Erodium</i> spp.	All States	350 mL/ha plus Agral in 200 L water	1 day grazing	Early autumn application.	Heavy grazing is necessary to reduce lucerne to 2cm in height before spraying.
			700 mL/ha plus Agral in 200 L water		Late winter application	
Oil seed poppies	Weed control	Tas only	300 mL to 1.5 L/ha	1 day grazing	Use in accordance with recommendations made by Department of Primary Industries of the poppy contracting company. <b>DO NOT add Agral or any other wetting agent to the spray solution.</b>	



**General weed control (continued):**

Crop	Weeds Controlled	State	Rate *	WHP	Critical Comments	
Orchards and Vineyards	Capeweed	All States	1.5 L plus 1.4 L Agral in 700 L water per ha plus 1.6 L/ha Paraquat 250.	Nil	Apply as a directed spray under trees or vines. Under most conditions, Paraquat 250 at 1.6 L/ha will give effective control of grasses and broadleaved weeds in orchards, but where heavy infestations of capeweed occur, Genfarm Diquat 200 should be added to Paraquat 250 at the rate of 1.2 L/ha. For inter-row or around butts, use high volume applications. For full ground cover use a minimum of 700mL/ha. Paraquat 250 and Genfarm Diquat 200 have no effect on brown bark but care should be taken when spraying around trees to avoid spray contacting green bark or plant material.	
Pasture Renovation and establishment	Capeweed and <i>Erodium</i> spp. (Storksbill)	All States	750 mL to 1.5 L/ha plus Agral in a minimum of 100 L water	1 day grazing	Apply by boom spray as an overall spray on 'run-down' pasture after heavy grazing. Pasture should not be greater than 4cm long when sprayed. Grazing should be carried out during previous spring, summer and early autumn. Where Capeweed is in the very young seedling stage (2 or 3 true leaves only) rates may be reduced to 350 mL/ha. Where Capeweed infestation is high, oversowing with new pasture seed by direct drilling is advisable. Direct drill 3 to 7 days after spraying using a pasture mixture suitable to the district.	
	Barley Grass, Brome Grass, Silver Grass and Sweet Vernal Grass		750 mL to 1.5 L/ha plus 1 to 2 L/ha Paraquat 250 in a minimum of 100 L water/ha			
Row crops, vegetables and market gardens	Broadleaf weeds	All States	1.4 L/ha	Nil	(seedling weeds)	Diquat/Paraquat 250 Herbicide and Paraquat 250 Herbicide are more generally used for grass and broadleaf weed control in these situations. However, where broadleaf weeds dominate, particularly capeweed, Genfarm Diquat 200 should be tank mixed with Paraquat 250 Herbicide or instead of Paraquat 250 Herbicide where grass weeds are absent. Apply as a blanket spray prior to crop emergence. Once crops have emerged, or seedlings have been transplanted, apply as a shielded spray between crop rows. Do not allow spray to contact any part of the crop.
			2.8 or 4 L/ha per 200 to 300 L water/ha plus 200 mL Agral per 100L water	Nil	(mature weeds)	
Wheat and Oats	Capeweed	Qld, NSW, Vic, Tas and SA only	550 mL/ha in 200 L of water	Nil	<b>Small seedlings. DO NOT</b> add wetting agent. Spray when the crop is between the 4 (wheat) or 3 (oats) leaf and early tillering stage.	

			700 mL/ha in 200 L of water	Nil	<b>Older seedlings: DO NOT</b> add wetting agent. Spray when the crop is between the 4 (wheat) or 3 (oats) leaf and early tillering stage.
--	--	--	--------------------------------------	-----	--

**General weed control (continued):**

Crop	Weeds Controlled	State	Rate *	WHP	Critical Comments	
Winter cereals	Pre-harvest weed control	All States	1 or 3 L/ha	Nil	Spray as soon as the crop is fully mature and ready for harvesting. Under wet spring conditions, crops can periodically become infested with weeds which seriously interfere with harvest operations.	
Wheat		NSW only	2 L/ha	Nil	Light to moderate stands.	Ensure that spray penetrates deep down into the crop canopy.
			3 L/ha	Nil	Moderate to heavy stands.	

**\*NOTE:** Use higher rate for dense or weedy crops.

WETTING AGENT: Add Agral at the rate of 200 mL/100 L or Wetter 1000 at 160 mL/100 L of prepared spray unless otherwise specified.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD:**

**Grazing:** DO NOT GRAZE OR CUT SPRAYED VEGETATION FOR STOCK FOOD FOR 1 DAY AFTER APPLICATION.

**Harvest:**

Cotton, dry beans, peas, mung beans,  
asparagus, hops, orchards and vineyards,  
row crops, vegetables and market gardens,  
oats, wheat and winter cereals:

**NOT REQUIRED WHEN USED AS DIRECTED**

Pigeon peas, canola, sunflower

soybeans, sugarcane:

**DO NOT HARVEST FOR 4 DAYS AFTER APPLICATION.**

Rice:

**DO NOT HARVEST FOR 5 DAYS AFTER APPLICATION.**

Potatoes:

**DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.**

Sweet Potatoes:

**DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.**

Poppies:

**DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION**

**DO NOT USE TREATED WATER FOR HUMAN CONSUMPTION, LIVESTOCK WATERING OR IRRIGATION PURPOSES FOR 10 DAYS AFTER APPLICATION.**

**GENERAL INSTRUCTIONS****USES**

Genfarm Diquat 200 is an aqueous solution of diquat, a non-volatile herbicide which unique properties. It very quickly kills green growth with which it comes into contact and is particularly effective against broadleaved weeds. It is inactivated on contact with the soil and crop roots and seeds below the soil remain unharmed. It can be safely applied around bushes and trees which have no green bark. It is non-volatile, easily mixed with water and active at low concentrations.

**RESISTANT WEEDS WARNING**

<b>GROUP</b>	<b>L</b>	<b>HERBICIDE</b>
--------------	----------	------------------

Genfarm Diquat 200 Herbicide is a member of the bipyridyl group of herbicides. The product has the inhibitor of photosynthesis at photosystem I mode of action. For weed resistance management, the product is a Group L herbicide.

Some naturally-occurring weed biotypes resistant to the product and other inhibitors of Group L herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. The resistant weeds will not be controlled by this product or other inhibitors of Group L herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Nutrien Ag Solutions Limited accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

### **MIXING**

Add the required quantity of Genfarm Diquat 200 to water in the spray tank and agitate to give even mixing. Agitate again if left standing. Use clean water only, as suspended soil particles in dirty water will interfere with herbicidal action.

### **WETTING AGENT**

Genfarm Diquat 200 Non-Residual Herbicide contains no wetting agent, and a non-ionic wetting agent must be added to the spray mixture unless otherwise specified. Add Agral at the rate of 200 mL/100 L or Wetter 1000 at 160 mL/100 L of prepared spray unless otherwise specified.

### **APPLICATION**

For best results, an even and complete coverage and good penetration of the spray into the target foliage is necessary. Best results will be obtained when application is made in dull weather or at the end of the day. Genfarm Diquat 200 is rapidly absorbed and is not affected by rain falling shortly after application.

### **APPLICATION RATES**

For application to seedling weeds, Genfarm Diquat 200 is generally recommended at 1.4 L/ha and Paraquat 250 Herbicide at 1.1 L/ha. Use Genfarm Diquat 200 at 2.8 to 4 L/ha and Paraquat 250 Herbicide at 2.2 to 3.2 L/ha when weeds are at the older stages of growth. Paraquat 250 Herbicide is preferred where grasses are dominant and Genfarm Diquat 200 where there are mainly broadleaf weeds.

Best results may be obtained when application is made in dull weather or at the end of the day. This product may also be used with 200 g/L Paraquat products at 1.4L/ha for seedling weeds and 2.8 to 4L/ha for older stage of weed growth.

### **BOOM SPRAYING**

A boom sprayer fitted with flat fan nozzles is preferred to ensure even coverage and to minimise drift. The boom should be set at sufficient height above the crop to provide a complete double overlap of the flat spray pattern. Spray drop arms on booms are useful for dense crops such as potatoes. A minimum spray volume of 100 L/ha is recommended.

### **HIGH VOLUME SPOT SPRAYING**

Hand-held equipment use 250 mL of product per 100 L of water and spray to visible wetness (about 700 to 1000 L/ha). Use 30 mL product plus 30 mL Agral per 15 litre knapsack.

### **AERIAL APPLICATION**

Flying height, pressure, nozzle size and positioning on the aircraft should be such as to minimise spray drift. Apply 30 to 60 L of spray per hectare. Avoid spraying in high winds or under temperature inversion conditions. Wash any spillage during filling of the aircraft and make sure there are no leaks in the spraying system. Inspect the aircraft regularly for signs of corrosion and ensure the paintwork is in good condition.

### **CAUTION – USE BY AIRCRAFT**

Although this product is no different in drift behaviour from other chemicals, it has a rapid spotting effect on green foliage and, as with all herbicides; special care must be taken to avoid drift onto adjacent crops. Aircraft operators must not apply during periods of thermal (temperature) instability, and should avoid wind conditions and flying heights conducive to drift.

### **WEED CONTROL IN ROW CROPS, VEGETABLES AND MARKET GARDENS**

#### **Pre-planting and pre-crop emergence:**

To control weeds in seed beds before sowing, or post-sowing pre-crop emergence, apply as a blanket spray with this product using boom spray equipment or knapsack sprayers.

#### **Post-emergence inter-row weed control:**

Use shielded nozzles for rapid control of weeds in inter-row spaces of row crops, after crop seedlings have emerged, or when transplanted crops are established. **Direct the spray so that it does not touch the crop.**

**Pre-harvest crop desiccation:**

Green crop foliage and weeds can seriously interfere with harvesting operations of a number of crops. This product can be used to facilitate harvesting by desiccating weeds, accelerating the drying of crops and reducing the moisture content of seeds. Drying costs are reduced, harvesting delays and associated risks avoided.

**WARNING**

**Markers** - if possible, fixed markers should be used. Human markers are not recommended unless flaggers are protected by engineering controls such as vehicles with enclosed cabs.

**COMPATIBILITY**

This product mixes readily with Paraquat 250 Herbicide, the soil residual herbicides Atrazine Granules, Diuron WG, Simazine Granules and other 200 g/L Paraquat products, where prolonged weed control is required as well as a quick knockdown.

**EXPORT OF TREATED PRODUCE**

Growers should note that MRL's or import tolerance do not exist in all markets for edible produce treated with Genfarm Diquat 200 Non-Residual Herbicide. If you are growing edible produce for export, please check with Nutrien Ag Solutions Limited for the latest information on MRL's and import tolerances BEFORE using Genfarm Diquat 200 Herbicide.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

**DO NOT** apply under weather conditions or from spraying equipment which may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

**PROTECTION OF LIVESTOCK**

Domestic pets and poultry – keep away from treated areas. Low hazard to bees. No special precautions are required.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

**DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Store in a locked room or a place away from children, animals, food, feedstuffs, seed and fertilisers.

This containers can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple or preferably pressure rinse containers before disposal. Dispose of rinsate by adding it to the spray tank. **DO NOT** dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection point or similar container management program site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

*For Refillable containers (110L, 1000L):* Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**SAFETY DIRECTIONS**

Very dangerous. Poisonous if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. **DO NOT** inhale spray mist. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves, face shield or goggles, and half-face respirator or disposable respirator.

If clothing becomes contaminated with product or wet with spray, remove clothing immediately. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles, respirator and if rubber wash with detergent and warm water, and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

**SAFETY DATA SHEET**

For further information refer to the Safety Data Sheet (SDS), which can be obtained from the supplier.

**CONDITIONS OF SALE**

The use of this product is beyond the control of Nutrien Ag Solutions Limited. Any provisions or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Nutrien Ag Solutions Limited does not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

APVMA Approval No: 64889/58177

Batch Number:

Date of Manufacture:

Nutrien Ag Solutions Limited

Level 5, Building A, 26 Talavera Road,

Macquarie Park, NSW 2113.

Tel: (02) 9889 5400

Product Support Tel: 1800 44 88 92

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Causes damage to organs through prolonged or repeated exposure; If inhaled remove person to fresh air and keep comfortable for breathing; If skin or eye irritation persists get medical attention; Use only outdoors or in a well-ventilated area; Contaminated work clothing should not be allowed out of the workplace.

<b>BIPYRIDILIUM PESTICIDES, LIQUID, TOXIC (CONTAINS DIQUAT)</b>	
<b>UN NO. 3016</b>	<b>PG III</b>
<b>HAZCHEM 2X</b>	
<b>IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE FOR SPECIALIST ADVICE IN AN EMERGENCY 1800 033 111 24 HOURS AUSTRALIA WIDE</b>	
