

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

GENFARM
AZOXY XTRA
FUNGICIDE

ACTIVE CONSTITUENTS: 200 g/L AZOXYSTROBIN
80 g/L CYPROCONAZOLE

GROUP **11** | **3** FUNGICIDE

Controls certain fungal diseases in Barley, Wheat and Peanuts
as specified in the Directions for Use.

IMPORTANT: READ THIS BOOKLET BEFORE USE

Nutrien Ag Solutions Limited
Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW, 2113
Tel: (02) 9889 5400 • Product Support Tel: 1800 44 88 92

DIRECTIONS FOR USE

RESTRAINTS

Cereals only:

DO NOT apply more than two applications of Genfarm Azoxy Xtra Fungicide (or any other strobilurin or Group 3 fungicide) in any one season on the same paddock.

DO NOT exceed a combined total of 800 mL/ha of Genfarm Azoxy Xtra Fungicide in any one season on the same paddock.

CROP	DISEASES	RATE/HA	CRITICAL COMMENTS
Barley	Leaf Rust (<i>Puccinia hordei</i>)	200-800 mL	<p>Apply from jointing (Z30) and before infection averages 10% on the flag-2 leaf. Ensure thorough coverage of stems and leaves. Use higher rates under high infection pressure or when longer residual protection is required. Lower rates are effective under low disease pressure but have reduced residual effect.</p> <p>Repeat application at early booting stage if necessary. Two applications of the lower rate will generally provide better control than a single application at the higher rate.</p> <p>Addition of Adigor[^] at 2% v/v improves disease control, especially at the lower rate.</p>
	Net Blotch: Net Form (<i>Pyrenophora teres f. teres</i>), Spot Form (<i>Pyrenophora teres f. maculata</i>)		
	Powdery Mildew (<i>Blumeria graminis hordei</i>)	400-800 mL	<p>Spray at the first sign of the disease during the tillering stage. A repeat spray 21 to 28 days later may be required. Ensure thorough coverage of stems and leaves. Higher rates provide longer protection.</p>

CROP	DISEASES	RATE/HA	CRITICAL COMMENTS
Peanuts	<p>Early Leaf Spot (<i>Cercospora arachidicola</i>), Late Leaf Spot (<i>Cercosporidium personatum</i>), Net Blotch (<i>Phoma arachidicola</i>), Rust (<i>Puccinia arachidicola</i>)</p>	500-750 mL	<p>Apply 1 to 2 applications of Genfarm Azoxy Xtra Fungicide at 10 to 14 day intervals from first flowering and before disease symptoms appear or when disease pressure is low. Use the low rate when disease pressure is low e.g. early season plantings, and use higher rates when weather conditions favour disease development and/or the crop has a large canopy. When targeting Net Blotch, or when weather conditions are not favourable for application, add a non-ionic surfactant at its recommended use rate.</p> <p>Resistance Management Disease control may be reduced if strains of pathogens less sensitive to Genfarm Azoxy Xtra Fungicide develop. DO NOT use Genfarm Azoxy Xtra Fungicide curatively. DO NOT apply more than 1/3 of the total number of fungicide applications per crop as Genfarm Azoxy Xtra Fungicide. For example, if the total number of fungicides applied to the crop is 9, DO NOT apply more than 3 applications of Genfarm Azoxy Xtra Fungicide. DO NOT exceed 3 applications of Genfarm Azoxy Xtra Fungicide per crop. DO NOT apply more than two consecutive applications of Genfarm Azoxy Xtra Fungicide. DO NOT apply more than 5 DMI sprays per season. Genfarm Azoxy Xtra Fungicide contains a DMI fungicide and should be included in the total number of DMI sprays.</p>

CROP	DISEASES	RATE/HA	CRITICAL COMMENTS
Wheat	Leaf Rust (<i>Puccinia triticina</i>)	400-800 mL	Apply after flag leaf is 70% emerged and before disease levels reach 1% of flag leaf area. Consider control if disease is greater than 5 to 10% on any lower leaf layer. Use higher rates under high infection pressure or when longer residual protection is required. Lower rates are effective under low disease pressure but have reduced residual effect. Ensure thorough coverage.
	Powdery Mildew (<i>Blumeria graminis tritici</i>)		Spray at the first sign of the disease during the tillering stage. A repeat spray 21 to 28 days later may be required. Ensure thorough coverage of stems and leaves. Higher rates provide longer protection.
	Stripe Rust (<i>Puccinia striiformis</i>)		Spray at the first sign of the disease and before 10 to 20% of leaves are infected. A repeat spray 21 to 28 days later may be required. Use higher rates under high infection pressure or when longer residual protection is required.
	Yellow Spot (Pyrenophora tritici-repentis)		Apply once between 70% flag leaf emergence and early flowering. Use higher rates under high infection pressure or where longer residual protection is required. Apply after 70% flag leaf emergence and before disease levels reach 5% on flag leaf. Higher rates give longer residual protection and often better economic returns.

WARNING

Azoxystrobin is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees.

DO NOT spray Genfarm Azoxy Xtra Fungicide where spray drift may reach apple trees.

DO NOT spray when conditions favour drift beyond the area intended for application. Conditions which may contribute to drift include thermal inversions, excessive wind speed, certain spray nozzle/pressure combinations, small spray droplet size, etc.

DO NOT use spray equipment which has been previously used to apply Genfarm Azoxy Xtra Fungicide to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity.

WITHHOLDING PERIODS

Barley, Wheat:

Grazing: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 21 DAYS AFTER APPLICATION.

Harvest: DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.

Peanuts:

Grazing: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 14 DAYS AFTER APPLICATION.

Harvest: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

MIXING

Genfarm Azoxy Xtra Fungicide is a suspension concentrate (SC) formulation that mixes readily with water and is applied as a spray.

1. Partly fill the spray tank with water.
2. Start the agitation.
3. Add the correct amount of product to the spray tank with the agitation system running.
4. Continue agitation while topping up the tank with water and while spraying.
5. Use the spray mix within 24 hours of preparation.

GROWING CONDITIONS

In general Genfarm Azoxy Xtra Fungicide should be applied under good growing conditions with adequate soil moisture. Genfarm Azoxy Xtra Fungicide should not be applied to crops that are under stress due to very dry, very wet, frosty conditions, nutrient deficiency or high insect pressure, as this may give less than reliable results.

SPRAY TIMING

Always inspect crops to assess disease development immediately before spraying. Best results will be achieved from applications made in the earliest stages of disease development or as a protectant treatment following a disease risk assessment.

Cereals: Applications of Genfarm Azoxy Xtra Fungicide should be made within the crop growth stages ranging from tillering to early flowering (Z26 to Z62). The object of spraying is to keep the upper 2 to 3 leaves green and functioning through grain filling stage.

Peanuts: The object of spraying is to control diseases and stop defoliation during pegging or pod fill stages to ensure stems are sufficiently strong for harvesting.

APPLICATION

May be applied by boom spray or aircraft. Ensure complete coverage of all leaves and stem is obtained.

Cereals

Boom application: As a guide, apply in spray volumes ranging from 50 to 100 L water per hectare. The use of flat fan nozzles is recommended using a fine to medium quality spray pattern, with pressures in the range 100 to 400 kPa.

Aerial application: As a guide, apply using 20 to 30 L water per hectare using a fine to medium quality spray. Use the lower rate when applications are made with a cross wind of not less than 5 knots. Use the higher rates when applying to dense crops.

Peanuts

Boom application: As a guide, apply in spray volumes ranging from 150 to 500 L water per hectare. The use of hollow cone nozzles or flat fans is recommended using a fine to medium quality spray pattern, with pressures in the range of 500 to 1000 kPa for the hollow cones and 150 to 300 kPa for flat fan nozzles.

Aerial application: As a guide, apply 25 to 60 L water per hectare using a fine to medium quality spray. Use with the lower rate when applications are made with a cross wind of not less than 5 knots. Use the higher rates when applying to dense crops.

COMPATIBILITY/TANK MIXING

Genfarm Azoxy Xtra Fungicide is physically compatible with a wide range of products, however the biological compatibility of these mixtures may not have been fully tested under all environmental and biological conditions.

A mixture of Genfarm Azoxy Xtra Fungicide with any other product may in some circumstances be ineffective or may cause damage.

If tank mixes are to be used observe all directions, precautions and limitations on all products to be used.

As formulations of other manufacturer's products are beyond the control of Landmark, and the quality of water may vary with location, all mixtures should be tested prior to mixing commercial quantities.

EXPORT SLAUGHTER INTERVAL

DO NOT slaughter animals destined for export within 7 days of consumption of treated cereal forage or straw, or peanut forage or fodder.

PRECAUTION

Re-entry Period: DO NOT allow entry into treated areas until the spray has dried unless wearing cotton overalls (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

FUNGICIDE RESISTANCE WARNING

Genfarm Azoxy Xtra Fungicide Fungicide is a combination of a DMI and a strobilurin and related products fungicide. For fungicide resistance management Genfarm Azoxy Xtra Fungicide is both a Group 11 and a Group 3 fungicide. Some naturally occurring individual fungi resistant to Genfarm Azoxy Xtra Fungicide and other Group 11 and/or Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Genfarm Azoxy Xtra Fungicide or other Group 11 and/or Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

GROUP	11	3	FUNGICIDE
-------	----	---	-----------

Since the occurrence of resistant fungi is difficult to detect prior to use, Nutrien Ag Solutions Limited accepts no liability for any losses that may result from the failure of Genfarm Azoxy Xtra Fungicide to control resistant fungi.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

HIGHLY TOXIC TO AQUATIC LIFE. DO NOT contaminate dams, waterways or drains with the chemical or used containers.

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift on adjacent areas, particularly wetlands, waterbodies or watercourses.

DO NOT apply Genfarm Azoxy Xtra Fungicide aerially if waterbodies, watercourses or wetlands are within 50 metres downwind of the application area.

STORAGE AND DISPOSAL

Store in closed original container in a cool, well-ventilated area as cool as possible. DO NOT store for prolonged periods in direct sunlight.

Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. Avoid contact with eyes. When opening the container and preparing spray wear: cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Poisons Information Centre Phone Australia 131126.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Nutrien Ag Solutions Limited. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Nutrien Ag Solutions Limited accepts no responsibility for any consequences resulting from the use of this product. Nutrien Ag Solutions Limited will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.



Nutrien Ag Solutions Limited

Level 5, Building A, 26 Talavera Road,
Macquarie Park, NSW, 2113 • Tel: (02) 9889 5400

Product Support Tel: 1800 44 88 92