

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**GENFARM**  
**IMAZA-COMBI**  
**HERBICIDE**

**ACTIVE CONSTITUENTS:**  
33 g/L IMAZAMOX present as the ammonium salt  
15 g/L IMAZAPYR present as the ammonium salt

**GROUP B HERBICIDE**

**For the early post-emergence control of certain annual grass and broadleaf weeds as part of the Clearfield<sup>^</sup> Production System for Clearfield Plus wheat, Clearfield barley, and Clearfield canola, as specified in the DIRECTIONS FOR USE section of this label**

**IMPORTANT: READ THIS BOOKLET BEFORE USE**

**Nutrien Ag Solutions Limited**  
**Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW, 2113**  
**Tel: (02) 9889 5400 • Product Support Tel: 1800 44 88 92**

**DIRECTIONS FOR USE**

**RESTRAINTS**

Apply ONLY to certified Clearfield Plus wheat, Clearfield barley and Clearfield canola varieties with the Clearfield technology, excluding Clearfield STL and Clearfield JNZ. DO NOT apply to conventional or other herbicide tolerant canola, wheat, and barley varieties.

DO NOT apply to crops that are stressed due to conditions such as waterlogging, too little moisture, frost, disease or nutritional disorders.

DO NOT apply by aircraft.

DO NOT apply if rain is expected within 2 hours of application.

DO NOT apply more than once per season to any one crop.

DO NOT use in Clearfield Plus wheat crops in tank mix or sequentially with diuron, sulfonyleureas, or sulfonamides.

**SPRAY DRIFT RESTRAINTS**

DO NOT apply with spray droplets smaller than a MEDIUM spray droplet size category according to nozzle manufacturer specifications that refer to the ASAE S572 Standard or the BCPC Guideline.

DO NOT apply when wind speed is less than 3 or more than 20 kilometers per hour as measured at the application site.

DO NOT apply during surface temperature inversion conditions at the application site.

Users of this product MUST make an accurate written record of the details of each spray application within 24 hours following application and KEEP this record for a minimum of 2 years.

The spray application details that must be recorded are: 1 date with start and finish times of application; 2 location address and paddock/s sprayed; 3 full name of this product; 4 amount of product used per hectare and number of hectares applied to; 5 crop/situation and weed/pest; 6 wind speed and direction during application; 7 air temperature and relative humidity during application; 8 nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application; 9 name and address of person applying this product. (Additional record details may be required by the state or territory where this product is used.)

**MANDATORY NO-SPRAY ZONES**

DO NOT apply if there are aquatic or wetland areas including aquacultural ponds within 50 meters downwind from the application area.

DO NOT apply if there are non-target vegetation or animal habitat within 5 meters downwind from the application area.

CROP USE OR SITUATION	WEEDS CONTROLLED	RATE
<p><b>Clearfield Plus wheat</b></p> <p>DO NOT use on CL STL and CL JNZ wheat varieties.</p> <p><b>Clearfield barley</b></p>	<p>Brome (<i>Bromus diandrus</i> and <i>B. rigidus</i>), Barley (<i>Hordeum vulgare</i>), Barley grass (<i>Hordeum leporinum</i>), Indian hedge mustard (<i>Sisymbrium orientale</i>), Muskweed (<i>Myagrum perfoliatum</i>), Oat (<i>Avena sativa</i>), Triticale (<i>Triticosecale</i> spp.), Wheat (<i>Triticum aestivum</i>) – non <b>Clearfield</b> varieties, Wild oat (<i>Avena fatua</i>), Wild radish (<i>Raphanus raphanistrum</i>), Wild turnip (<i>Brassica tournefortii</i>)</p>	<p>375-750 mL/ha plus Hasten<sup>^</sup> or Kwickin<sup>^</sup> (or equivalent) at 0.5 L/100 L spray volume</p>
	<p>Annual ryegrass (<i>Lolium rigidum</i>), Dense flower fumitory (<i>Fumaria densiflora</i>), Marshmallow (<i>Malva parviflora</i>), Sub clover (<i>Trifolium subterraneum</i>),</p> <p><b>Suppression</b></p> <p>Bedstraw species. (<i>Galium tricornutum</i> and <i>G. aparine</i>), Doublegee (<i>Emex australis</i>), Silver grasses (<i>Vulpia bromoides</i> and <i>V. myuros</i>)</p>	<p>600-750 mL/ha plus Hasten or Kwickin (or equivalent) at 0.5 L/100L spray volume</p>
<p>Canola varieties with <b>Clearfield</b> technology <b>only</b></p>	<p>Indian hedge mustard (<i>Sisymbrium orientale</i>), Muskweed (<i>Myagrum perfoliatum</i>), Wild radish (<i>Raphanus raphanistrum</i>), Wild turnip (<i>Brassica tournefortii</i>)</p>	<p>300–500 mL/ha plus Hasten or Kwickin (or equivalent) at 0.5 L/100L spray volume</p>
	<p>As above plus:</p> <p>Capeweed (<i>Arctotheca calendula</i>), Field pea (<i>Pisum sativum</i>), Narrow leaf lupin (<i>Lupinus angustifolius</i>), Sub clover (<i>Trifolium subterraneum</i>)</p>	<p>300-500 mL/ha plus Clopyralid 600 Herbicide at 75 to 150 mL/ha plus Hasten or Kwickin (or equivalent) at 0.5 L/100L spray volume</p>

CRITICAL COMMENTS
<p>NOT for use on <b>Clearfield</b> wheat varieties <b>Clearfield</b> STL and <b>Clearfield</b> JNZ.</p> <p>Apply to Clearfield Plus wheat crops from the 3 leaf stage to 1st node stage (Z31). Apply to Clearfield barley crops from the 5 leaf stage to 1st node stage (Z31).</p> <p>DO NOT apply to Clearfield Plus wheat or Clearfield barley after the 1st node stage (Z31).</p> <p>Applications should be targeted at grass weeds when the majority are in the 2-4 leaf stage and only when within the recommended crop stages.</p> <p>Application to multi-tillered crops may impair weed control because of poor contact and coverage of weeds.</p> <p>– see Compatibility. Tank mixes with Clopyralid 600 Herbicide at 75 mL/ha will provide control of composite and legume weeds. Tank mixes with MCPA 570 LVE Herbicide at 500 mL/ha will provide control of composite and brassicaceous weeds. The control of brassicaceous weeds will depend on the status of Group B resistance in the population. The addition of MCPA 570 LVE will improve control and provide an additional mode of action for resistance management. If other weeds require control, apply appropriate herbicides at least two weeks before or after Imaza-Combi and only when signs of regrowth or renewed vigor appear, otherwise the effects of the early treatment may affect the performance of the subsequent treatment.</p>
<p>Weed species will either be controlled or suppressed. In both cases, surviving plants will be stunted and will be uncompetitive with the crop, and seed set will be prevented or greatly reduced.</p> <p>The control of annual ryegrass varies from excellent to poor depending on the status of Group B resistance in the population and environmental conditions. Where the population is expected to exceed 200 plants/m<sup>2</sup> or a high level of control is required, or the ryegrass is known to be resistant or thought to be developing resistance, an application of a suitable pre-emergent should be made prior to sowing. A follow up grass selective herbicide may also be necessary.</p>
<p><b>Read Follow Crop comments and restrictions on the label prior to use.</b></p> <p>Apply to crop at the 2 to 6 leaf stage.</p> <p>DO NOT apply after the 6 leaf stage.</p> <p>Apply to actively growing weeds in the 3-leaf to 2-tiller stage and broadleaf weeds in the 2 to 6 leaf stage.</p> <p>Use the higher rate when weed numbers are high or towards the upper end of the recommended growth stages, or when the crop is at the 5 to 6 leaf stage to ensure better contact and coverage.</p> <p>If other weeds require control, apply appropriate herbicides at least two weeks after <b>Imaza-Combi</b> and only when signs of regrowth or renewed vigour appear, or the effects of <b>Imaza-Combi</b> may affect their performance.</p>
<p>Refer to critical comments for 300 to 500 mL/ha alone.</p> <p>Clopyralid 600 Herbicide aids in the control of legume and composite weed species. Refer to the Clopyralid 600 Herbicide label. Use rates above 75 mL/ha when these weeds are primary weeds in the paddock and when required by their growth stage.</p> <p>Clopyralid 600 Herbicide above 75 mL/ha can slightly impair grass control. The addition of Clopyralid 600 Herbicide does not affect the control of wild radish and wild turnip.</p> <p>(Refer to the Compatibility section of this label and the Clopyralid 600 Herbicide label for further details of use).</p>

CROP USE OR SITUATION	WEEDS CONTROLLED	RATE
Canola varieties with <b>Clearfield</b> technology only – <i>continued</i>	Non- <b>Clearfield</b> Barley ( <i>Hordeum vulgare</i> ), Barley grass ( <i>Hordeum leporinum</i> ), Great brome ( <i>Bromus diandrus</i> ), Indian hedge mustard ( <i>Sisymbrium orientale</i> ), oat ( <i>Avena sativa</i> ), Rigid brome ( <i>Bromus rigidus</i> ), Marshmallow ( <i>Malva parviflora</i> ), Sub clover ( <i>Trifolium subterraneum</i> ), Non- <b>Clearfield</b> wheat ( <i>Triticum aestivum</i> ), Wild oat ( <i>Avena fatua</i> ), Wild radish ( <i>Raphanus raphanistrum</i> ), Wild turnip ( <i>Brassica tournefortii</i> )  <b>Suppression only:</b> Annual ryegrass ( <i>Lolium rigidum</i> ), Bedstraw species ( <i>Galium tricornutum</i> and <i>G. aparine</i> ), Doublegee ( <i>Emex australis</i> ), Silver grasses ( <i>Vulpia bromoides</i> and <i>B. myuros</i> )	600-750 mL/ha plus Hasten or Kwickin (or equivalent) at 0.5 L/100L spray volume
	As above plus: Capeweed ( <i>Arctotheca calendula</i> ), Field pea ( <i>Pisum sativum</i> ), Narrow leaf lupin ( <i>Lupinus angustifolius</i> )	600-750 mL/ha plus Clopyralid 600 Herbicide at 75-150 mL/ha plus Hasten or Kwickin (or equivalent) at 0.5 L/100L spray volume

CRITICAL COMMENTS
Apply to crop at the 2 to 6 leaf stage. DO NOT apply after the 6 leaf stage.  Apply to actively growing grass weeds in the 3-leaf to 2-tiller stage and broadleaf weeds in the 2 to 6 leaf stage.  Use the higher rate when weed numbers are high or towards the upper end of the recommended growth stages, or when the crop is at the 5 to 6 leaf stage, to ensure better contact and coverage.  Weeds will either be killed in high numbers (weeds controlled) or in moderate numbers (weeds suppressed). In both cases, surviving plants will be stunted and will be uncompetitive with the crop, and seed set will be prevented or greatly reduced.  If other weed species require control, apply appropriate herbicides at least two weeks after Imaza-Combi and only when signs of regrowth or renewed vigour appear or the effects of Imaza-Combi may affect their performance.  3 The control of annual ryegrass varies from excellent to poor depending on the status of Group B resistance in the population and environmental conditions. Where the population is expected to exceed 200 plants per sqm, or a high level of control is required, or the ryegrass is known to be resistant or thought to be developing resistance, an application of a pre-emergent herbicide should be made prior to sowing. A tank mix with a grass selective herbicide may also be necessary.
Refer to critical comments for 600 to 750 mL/ha alone.  DO NOT apply after the 6 leaf stage.  Clopyralid 600 aids in the control of legume and composite weed species. Refer to the Clopyralid 600 label. Use rates above 75 mL/ha when these weeds are primary weeds in the paddock and when required by their growth stage.  Clopyralid 600 above 75 mL/ha can slightly impair grass control.  The addition of Clopyralid 600 does not affect the control of other weeds controlled by Imaza-Combi.  (Refer to the Compatibility section of this label and the Clopyralid 600 label)

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

#### WITHHOLDING PERIOD

#### GRAZING

**CLEARFIELD PLUS WHEAT AND CLEARFIELD BARLEY: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.**

**CLEARFIELD CANOLA: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 5 WEEKS AFTER APPLICATION.**

#### HARVEST

**ALL CROPS: NOT REQUIRED WHEN USED AS DIRECTED.**

## GENERAL INSTRUCTIONS:

Imaza-Combi is for use in Clearfield Plus wheat, Clearfield barley, and Clearfield canola. Varieties with Clearfield technology are those that have been bred specifically to be tolerant to Imaza-Combi.

Imaza-Combi is absorbed through the leaves, green stems and roots of susceptible weeds and moved from the point of contact throughout the plant. Weeds will either die or will remain stunted and will not compete with the crop. Symptoms of kill may take one to two weeks to develop with death occurring up to one month from treatment. Symptoms first appear at the growth points where young foliage becomes discoloured and distorted before dying.

Imaza-Combi is primarily a post-emergence product. Best results will be achieved when good contact and coverage of weeds occurs and weeds are actively growing. The product must be mixed with Hasten or Kwickin (or equivalent) as per the Directions for Use.

Imaza-Combi also has some residual soil activity under good soil moisture conditions. Residual effects on weeds can be reduced when dry soil conditions follow application before the herbicide has moved to the root zone. Best results will be achieved when application is made to moist soil or if approximately 10 mm rain follows within several days of application.

Vigorous crop growth will assist in suppressing weeds not completely killed and those germinating later.

## MIXING

Imaza-Combi is a soluble concentrate formulation. Pour the required amount of the product into a spray tank containing almost the total amount of water required. Mix thoroughly. If Imaza-Combi is added during filling, foaming may occur. If excessive foaming becomes a problem, add a silicone based antifoaming agent at the manufacturers recommended rate. Do NOT use a suction probe unless the antifoaming agent has already been added to the spray tank water. Consult your distributor for specific information on suitable antifoaming agents.

When tank mixing this product with other recommended compatible products, first add the other product(s) to the tank and mix thoroughly before adding Imaza-Combi.

Imaza-Combi may be applied in hard or soft water.

The product is corrosive to mild steel. Use ONLY stainless steel, fibreglass, plastic or plastic-lined containers for mixing, storage and application.

## APPLICATION

DO NOT apply by aircraft.

Apply in minimum 70L water per hectare. When the crop is very leafy or when the total weed population exceeds 200plants/m<sup>2</sup>, apply in a minimum of 100L water per hectare to improve contact and coverage. Imaza-Combi should be applied a minimum of two hours before rainfall or irrigation. If tank-mixed with other products, follow recommendations for the mixing partner should these extend beyond two hours.

## EQUIPMENT CLEAN-UP

Thoroughly flush all spray equipment with water following the use of Imaza-Combi and before use with other products. If tank-mixed with other products, also follow clean-up procedures recommended for the mixing partner.

## COMPATIBILITY

Imaza-Combi is chemically compatible with the insecticide alpha-cypermethrin. It is also chemically and biologically compatible with the herbicides bromoxynil, bromoxynil + MCPA, clopyralid 300 and 600, MCPA LVE, MCPA DMA and tralkoxydim.

DO NOT tank mix with foliar fertilisers.

Mixes with clopyralid in Clearfield Plus wheat and Clearfield barley:

Tank mixes with clopyralid 600 at 150mL/ha will provide control of composite and legume weeds. Refer to clopyralid label.

Mixes with clopyralid in Clearfield canola:

Clopyralid aids in the control of legume and composite weed species, such as Annual Medics (*Medicago* spp.), Capeweed (*Arctotheca calendula*), Chickpea (*Cicer arietinum*), Faba bean (*Vicia faba*), Field pea (*Pisum sativum*), Lentil (*Lens culinaris*), Narrow leaf lupin (*Lupinus angustifolius*). Use rates above 150mL/ha when these weeds are primary weeds in the paddock and when required by their growth stage. C above 150mL/ha can slightly impair grass control. For Chickpea, Faba bean, Lentil: If targeting Chickpeas and Lentils up to 6 leaf stage and Faba beans up to 4 leaf stage, use a tank mix of Imaza-Combi with 250mL/ha clopyralid 600. The addition of clopyralid 600 does not affect the control of other weeds controlled by Imaza-Combi. Refer to the clopyralid 600 label.

## RESISTANT WEEDS WARNING

GROUP	B	HERBICIDE
-------	---	-----------

Genfarm Imaza-Combi Herbicide is a member of the imidazolinone group of herbicides. The product has the inhibitors of acetolactate-synthase (ALS) mode of action. For weed resistance management, the product is a Group B herbicide.

Some naturally-occurring weed biotypes resistant to the product and other Group B herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group B herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nutrien Ag Solutions Limited accepts no liability for any losses that may result from failure of this product to control resistant weeds.

## PRECAUTIONS

### RE-ENTRY

DO NOT re-enter treated areas until spray has dried. If re-entry is necessary wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves.

## BEST MANAGEMENT PRACTICE (BMP) PROGRAM

A detailed program has been developed that outlines sound agronomic and integrated weed management practices, designed to optimise the performance of the Clearfield Production Systems and minimise the potential for the development of herbicide resistance in weed populations.

Consultation on BMP should be undertaken with a Clearfield-accredited agronomist prior to use of Imaza-Combi in the Clearfield Production Systems for wheat, barley and canola. Implementation of the BMP is an essential part of herbicide resistance management.

## FOLLOWING CROPS

This product is broken down in the soil by microbes in wet, aerobic conditions. Under conditions that do not favour breakdown, carry-over soil residues can affect susceptible follow crops. Normally safe residue levels may still affect follow crops when soil nutrition is low or marginal, when cold and very wet soil conditions prevail, or when soil pathogens or nematodes are present. As environmental and agronomic factors make it impossible to eliminate all risks associated with this product, rotational crop injury is always possible.

Note: when the intention is to grow cereals on Clearfield canola stubble (treated with Imaza-Combi) self-sown canola volunteers must be removed before they mature beyond 2-leaf, all macro and micro-nutrients must be maintained at levels necessary to grow the planned crops, and sulfonylureas must not be used.

The following minimum re-cropping intervals (months after application) should be observed.

Months after Application	Follow Crops
0	Clearfield Plus wheat, Clearfield wheat, Clearfield barley, Clearfield canola
10	Chickpeas, Faba beans, Field peas, Lucerne, Lupins, Pasture legumes, vetch, Oats*, Triticale*, Non-Clearfield Barley, Non-Clearfield Wheat
34	Conventional and other herbicide tolerant canola, All other crops

\* Non-Clearfield Barley, Oats, Triticale, Non-Clearfield Wheat:

The following additional requirements apply if it is intended to sow these cereals during the next winter season:

- DO NOT apply Imaza-Combi later than the end of August (no later than the end of July in WA).
- DO NOT use Imaza-Combi in areas where rainfall from spraying to sowing of cereals is expected to be below 150 mm (for 300 – 375 mL/ha use), 200 mm (for up to 500 mL/ha use) and 250 mm (for 600 – 750 mL/ha use).
- DO NOT use above 375 mL/ha in the Lower Great Southern region of Western Australia.

For all situations, if expected rainfall is not received following use of Imaza-Combi, consult your local Nutrien Ag Solutions representative before planting non-Clearfield cereals. In calculating rainfall actually received, place greater emphasis on rain received from application up to the end of Spring and lesser emphasis on break rains. If single isolated heavy summer and autumn falls and break rains are required to achieve rainfall targets, it may not be safe to sow non-Clearfield cereals within 10 months of application. Consult your local Nutrien Ag Solutions representative for advice.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT spray within 50 m of wetlands or waterways.

## CROP SAFETY

This product may, in some circumstances, lead to transient crop yellowing and temporary slowing of growth of Clearfield Plus wheat, Clearfield barley and Clearfield canola but plants soon recover and yield is unaffected. This effect may be more pronounced when the product is used under poor growth conditions.

The Clearfield wheat varieties Clearfield STL and Clearfield JNZ have limited tolerance to Imaza-Combi. Application of Imaza-Combi to these wheat varieties can cause unacceptable injury especially under cold, wet conditions. Imaza-Combi should not be used on these varieties.

DO NOT use this product on any canola variety other than certified varieties with Clearfield technology. Extreme crop damage and/or death will result to conventional and other herbicide tolerant wheat and canola varieties.

## PROTECTION OF LIVESTOCK

Imaza-Combi is of low hazard to bees.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

### Refillable containers

Empty containers fully into application equipment. Close all valves and return to point of supply for refill or storage.

## **SAFETY DIRECTIONS**

Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the containers and preparing spray, wear cotton overalls (or equivalent clothing) buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing. When tank mixing with other products consult also the safety directions for those products.

## **FIRST AID**

If poisoning occurs, contact a Poisons Information Centre Phone Australia 131126. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, wash out immediately with water.

## **SAFETY DATA SHEET**

For further information refer to the Safety Data Sheet, which can be obtained from the supplier.

## **CONDITIONS OF SALE**

The use of this product is beyond the control of Nutrien Ag Solutions Limited. Any provisions or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Nutrien Ag Solutions Limited does not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.



## **Nutrien Ag Solutions Limited**

Level 5, Building A, 26 Talavera Road,  
Macquarie Park, NSW, 2113 • Tel: (02) 9889 5400

**Product Support Tel: 1800 44 88 92**