

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

GENFARM
PROPYZAMIDE 500SC
HERBICIDE

ACTIVE CONSTITUENT: 500 g/L PROPYZAMIDE

GROUP	D	HERBICIDE
-------	----------	-----------

For the selective control of certain grasses and broadleaf weeds in lettuce and legume seed crops and pastures as per the Directions for Use table.

IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING THIS PRODUCT

Nutrien Ag Solutions Limited
Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW, 2113
Tel: (02) 9889 5400 • Product Support Tel: 1800 44 88 92

DIRECTIONS FOR USE

CROP	WEEDS CONTROLLED	STATE	RATE (L/ha)
Lettuce Direct sown	Barleygrass, Blackberry nightshade Chickweed, Portulaca, Winter grass,	All States	4.5 in 330-550 L of water
Lettuce Transplanted	Rye grass Summer grass, Prince of Wales feather Nettles, Shepherds purse Wireweed	Vic only	2.3 L in 330- 550 L water
Lucerne, Clover, Medics etc grown for forage, hay or seed production	Annual ryegrass, Winter grass, Barley grass, Silver grass, Prairie grass, Canary grass, Great Brome, Squirrel Tail fescue and Rats Tail fescue	NSW, ACT, SA, WA, Tas only	1 – 1.5 L/ha
	Wild Oats, Perennial Ryegrass, Bent Grass, English Couch, Yorkshire Fog Grass, Chickweed, Seedling Sorrel, Blackberry Nightshade, Patterson's Curse, Seedling Wireweed		1.5 – 2 L/ha

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHTHOLDING PERIOD: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION

GENERAL INSTRUCTIONS

Best results are obtained by applying product to cool, wet soil.

Mixing

Prior to pouring, shake container vigorously, then add the required quantity of Genfarm Propyzamide 500SC to water in the spray vat while stirring or with agitators in motion. DO NOT mix with other pesticides.

RESISTANT WEEDS WARNING

GROUP	D	HERBICIDE
-------	---	-----------

Genfarm Propyzamide 500SC Herbicide is a member of the Benzamide group of herbicides. Genfarm Propyzamide 500SC Herbicide has the inhibitors of microtubule assembly mode of action. For weed resistance management, the product is a Group D herbicide.

Some naturally occurring weed biotypes resistant to Genfarm Propyzamide 500SC Herbicide and other Group D herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Genfarm Propyzamide 500SC Herbicide or other Group D herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Nutrien Ag Solutions Limited accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

Advice as to strategies and alternative and alternative treatments that can be sued can be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or Nutrien Ag Solutions representative.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

PROTECTION OF LIVESTOCK

DO NOT graze sheep on treated areas or feed to poultry or pigs.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

CRITICAL COMMENTS
Apply a pre-emergent immediately after sowing. To ensure best results, prepare the seed-bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm should be applied within 24 hours.
Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm should be applied within 24 hours.
SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed. ESTABLISHED CROPS: Apply any time before end of July. WEEDS: Apply when weeds are at post or Pre-emergent stages. BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25 mm of rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers and product.

Refillable containers (110L, 1000L only)

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length PVC gloves.

After use wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor, or Poisons Information Centre. Phone Australia: 131126.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Nutrien Ag Solutions Limited. Any provisions or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Nutrien Ag Solutions Limited does not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

APVMA Approval No: 66867/118811



Nutrien Ag Solutions Limited

Level 5, Building A, 26 Talavera Road,
Macquarie Park, NSW, 2113 • Tel: (02) 9889 5400

Product Support Tel: 1800 44 88 92