

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

GENFARM TEBU T

FLOWABLE FUNGICIDE/INSECTICIDE SEED TREATMENT

ACTIVE CONSTITUENTS: 25 g/L TEBUCONAZOLE
4 g/L TRIFLUMURON

GROUP **3** FUNGICIDE

GROUP **15A** INSECTICIDE

Wheat: For control of bunt, flag smut and loose smut.
Barley: For control of covered and loose smut of barley.
Oats: For control of covered and loose smut.
For protection against insect pests of stored seed grain.

IMPORTANT: READ THIS BOOKLET BEFORE USE

Nutrien Ag Solutions Limited
Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW, 2113
Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92

DIRECTIONS FOR USE

Restraint: DO NOT treat physically damaged seed.

CROP	DISEASE/PEST	RATE	CRITICAL COMMENTS
Wheat	Bunt, Flag Smut (seed and soil borne), Loose Smut	100 mL/ 100 kg seed	Genfarm Tebu T Flowable Fungicide/ Insecticide Seed Treatment should be mixed with water to give even coverage of seed.
Barley	Covered and Loose Smut		The quantity of water used for mixing will vary depending on type of equipment and quality of seed.
Oats	Covered and Loose Smut		Use a minimum of 400 mL of mixture (i.e. Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment plus water) with each 100 kg of seed.
Wheat, Barley, Oats	Protection against pests of stored grains: Granary Weevil, Lesser Grain Borer, Rice Weevil, Rust-red Flour Beetle, Sawtoothed Grain Beetle		DO NOT use more than 600 mL of mixture (product + water) with each 100 kg of seed. Whatever dilution is used it is essential that 100 mL of Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment is applied/100 kg seed. The mixture should be gently stirred regularly. If nasal irritation occurs wear a face mask while treating seed.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

WITHHOLDING PERIODS:

FOOD PRODUCING CROPS: NOT REQUIRED WHEN USED AS DIRECTED.

**STOCK: DO NOT GRAZE PLANTS GROWN FROM TREATED SEED, OR CUT FOR STOCK
FOOD WITHIN 4 WEEKS OF SOWING.**

GENERAL INSTRUCTIONS

In Wheat: Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment, may delay emergence. Crop will recover rapidly in moist soil types. However, treated seed should not be sown in soils that are likely to clod or crust over after sowing. Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment treated wheat deeper than 50 mm may slightly reduce percentage plant establishment in heavier soils and sowing rates may need to be adjusted accordingly. Germination may be reduced if poor quality seed is used (cracked or shrivelled) – best to use high quality seed. Treated seed may be held over for sowing in the following season. If so, additional insecticide treatment may be necessary. If the seed is not used immediately after treatment, it should be stored in a dry, well-ventilated place. Treated seed which will not be used for sowing at all should be destroyed.

FUNGICIDE RESISTANCE WARNING

Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide.

GROUP 3 FUNGICIDE

Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Nutrien Ag Solutions Limited accepts no liability for any losses that result from failure of this product to control resistant fungi.

INSECTICIDE RESISTANCE WARNING

For insecticide resistance management Genfarm Tebu T Flowable Fungicide/Insecticide Seed

GROUP 15A INSECTICIDE

Treatment is a Group 15A insecticide. Some naturally occurring insect biotypes resistant to Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment and other Group 15A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment or other Group 15A insecticides are used repeatedly. The effectiveness of Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Nutrien Ag Solutions Limited accepts no liability for any losses that may result from the failure of Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment to control resistant insects.

Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment may be subject to specific resistance management strategies. For further information contact your local supplier, Nutrien Ag Solutions Limited representative or local agricultural department agronomist.

MIXING for use in a pickle pot or similar (liquid application)

The required quantity of Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment should be thoroughly mixed with a small quantity of clean, filtered water to make a creamy mix. Add this blend to the required quantity of water in the tank and mix thoroughly. Agitate prior to and during spraying.

DO NOT mix more than can be used for the day's seeding.

DO NOT tank mix with trace elements or other chemicals without first check compatibility.

SEED QUALITY

Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment should not be used on seed with more than 12% moisture content, as the treatment can raise the level by 0.6%. Seed to be treated should be of the highest quality. Poor quality (sprouted, cracked, pinched etc.) seed should not be used.

CLEANING OF EQUIPMENT

Clean equipment (augers etc.) with water after use. If changing from this water based product to a solvent based formulated product or vice versa, ensure complete dryness with Methylated spirits (or similar). Failure to do so will result in clogging of equipment.

PRECAUTION

DO NOT allow seed treated with this product to contaminate grain intended for human consumption.

If possible, use a separate auger for treating seed. If using the same auger for treating grain and moving grain for human consumption, ensure auger is completely cleaned as above.

PROTECTION OF LIVESTOCK

Seed treated with this product must not be used for animal consumption, poultry feed or mixed with animal feed.

DO NOT allow seed treated with this product to contaminate grain intended for animal consumption.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed treated seed or otherwise expose to wild or domestic birds or other animals.

Harmful to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Used containers or bags which have held treated seed must be correctly disposed of.

STORAGE AND DISPOSAL

When treated seed is stored it should be kept apart from other grain. Such bags or other containers should clearly be marked to indicate the contents have been treated with Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment.

DO NOT use seed treated with Genfarm Tebu T Flowable Fungicide/Insecticide Seed Treatment for human consumption. Bags which have held seed treated with this product are not to be used for any other purpose.

For Non-Refillable Containers: Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple or preferably pressure rinse containers before disposal. Dispose of rinsate by adding it to the spray tank. DO NOT dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection point or similar container management program site. The cap should not be replaced but may be taken separately. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers and product.

For Refillable Containers: Empty contents fully into application equipment. Close all valves and return to point of supply.

SAFETY DIRECTIONS

Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Nutrien Ag Solutions Limited. Any provisions or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Nutrien Ag Solutions Limited does not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.



Nutrien Ag Solutions Limited

Level 5, Building A, 26 Talavera Road,
Macquarie Park, NSW, 2113 • Tel: (02) 9889 5400

Product Support Tel: 1800 44 88 92