

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Genfarm

Pendimethalin 440 EC

HERBICIDE

ACTIVE CONSTITUENT: 440 g/L PENDIMETHALIN
SOLVENT: 501 g/L LIQUID HYDROCARBON

GROUP 3 HERBICIDE

For the control of wireweed, annual grasses (including ryegrass) and certain broadleaved weeds in various crops as shown in the DIRECTIONS FOR USE table.

IMPORTANT: READ THIS LEAFLET BEFORE USE

APVMA Approval No: 91417/132175

Nutrien Ag Solutions Limited
Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW, 2113
Tel: (02) 9889 5400 • Product Support Tel: 1800 44 88 92

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT sow sensitive summer crops such as sorghum and millets within 12 months of application of Genfarm Pendimethalin 440 EC Herbicide.
DO NOT apply to a poorly prepared seedbed.
DO NOT use Genfarm Pendimethalin 440 EC Herbicide mixed with atrazine on heavy clay soils.
DO NOT apply Genfarm Pendimethalin 440 EC Herbicide where waterlogging is likely to occur.

Pre-plant Application (Incorporated by the Sowing Process):		
SITUATION / CROP	WEEDS CONTROLLED	STATE
Barley and wheat (Including crops with undersown Lucerne)	Annual Ryegrass (<i>Lolium rigidum</i>), Wireweed (<i>Polygonum aviculare</i>), Suppression of Wild Oats (<i>Avena</i> spp.)	All States
Canola (Including crops with undersown Lucerne)	Suppression only of Silver Grass (<i>Vulpia</i> spp.)	Southern NSW, ACT, Vic, SA, WA, Tas only
Chickpeas, Faba Beans, Field Peas, Lentils, Lupins and Safflower	Annual Ryegrass (<i>Lolium rigidum</i>), Wireweed (<i>Polygonum aviculare</i>), Suppression only of Silver Grass (<i>Vulpia</i> spp.), Suppression of Wild Oats (<i>Avena</i> spp.)	Southern NSW, ACT, Vic, SA, WA, only
Lupins		WA only
		NSW, ACT, Vic, SA only

RATE / TREATED ha	CRITICAL COMMENTS
1.35 L/ha	<p>Barley and wheat (Including crops with undersown Lucerne): Where full (double) incorporation is impractical and where the seedbed tilth is fine and free of large stones and trash, apply Genfarm Pendimethalin 440 EC Herbicide up to 24 hours prior to sowing. Incorporate with the combine set to sow to a depth of 3-5cm. Use trailing harrows. (Refer to "Incorporation" under the heading General Instructions). In minimum till situations, only light incorporation will occur due to soil displaced by the points. Trailing harrows should be used to provide better incorporation and weed control. Avoid heavy trash situations which may result in uneven application and incorporation, leading to reduced weed control. Wheat and barley seed must be sown at least 1cm beneath the chemically treated band or reduced emergence may occur. Refer to "Cereal Seed Dressings" and "Weather Conditions at Planting" under GENERAL INSTRUCTIONS, for further information on emergence.</p> <p>Caution: Ensure thorough incorporation to maximise safety to undersown Lucerne.</p> <p>Canola (Including crops with undersown Lucerne): Apply Genfarm Pendimethalin 440 EC Herbicide up to 24 hours prior to sowing. Ensure Genfarm Pendimethalin 440 EC Herbicide is well incorporated. Sow Canola with a seeding depth of 1-3 cm, at the appropriate depth for conditions. Canola seed can be placed within the treated band. Incorporate with the combine set to sow the seed below the chemical band. DO NOT apply to Canola post plant pre-emergence. Use the higher rates on medium to heavy textured soils and the lower rates on light textured soils.</p> <p>Note: Development in waterlogged conditions after the application of Genfarm Pendimethalin 440 EC Herbicide may retard the emergence of Canola.</p> <p>If Wild Oats are expected to be a problem use a Wild Oat herbicide. Application to heavier soil types that have poor structural characteristics or are poorly worked may result in insufficient and uneven incorporation and unsatisfactory weed control. Heavy soils that are cloddy or have a surface crust or surface seal should be avoided. Attempting to incorporate Genfarm Pendimethalin 440 EC Herbicide when soils are very wet or very dry may also result in insufficient mixing and therefore poor weed control.</p> <p>Caution: Ensure thorough incorporation to maximise safety to undersown Lucerne. Use the lower rate when Canola is undersown with Lucerne.</p>
1.35 to 2.25 L/ha	
1.5 to 2.25 L/ha	
1.5 to 2.25 L/ha	<p>Where full (double) incorporation is impractical and where the seedbed tilth is fine and free of large stones and trash, apply Genfarm Pendimethalin 440 EC Herbicide or in the case of lupins Genfarm Pendimethalin 440 EC Herbicide plus Simazine up to 24 hours prior to sowing. Incorporate with the combine set to sow to a depth of 3-5cm. Use trailing harrows. (Refer to "Incorporation" under the heading General Instructions). Application to heavier soil types that have poor structural characteristics or are poorly worked may result in insufficient and uneven incorporation and unsatisfactory weed control. Heavy soils that are cloddy or have a surface crust or surface seal should be avoided. Attempting to incorporate Genfarm Pendimethalin 440 EC Herbicide when soils are very wet or very dry may also result in insufficient mixing and therefore poor weed control. Refer to "Weather Conditions at Planting" under GENERAL INSTRUCTIONS, for further information on emergence.</p> <p>If Wild Oats are expected to be a problem use a Wild Oat herbicide. Use the higher rates on medium to heavy textured soils and the lower rate on light textured soils.</p>
1.5L/ha plus simazine at 500g ai/ha	
1.5L/ha plus simazine at 1.5 kg ai/ha	

Pre-plant incorporation (Double Incorporation):

Note: For all crops listed in the double incorporation table below:

Incorporate Genfarm Pendimethalin 440 EC Herbicide into the soil within 24 hours of application. Refer to the notes on incorporation under the heading General Instructions. Where a range of application rates is provided, use the higher rate on heavy textured soils or those high in organic matter and the lower rate on light to medium textured soils. Application to soil types with poor structural type characteristics or that are poorly worked may result in insufficient or uneven incorporation and, subsequently less than satisfactory weed control. Heavy soils, which are cloddy or have a surface crust or surface seal, are to be avoided. Attempting to incorporate Genfarm Pendimethalin 440 EC Herbicide when soils are very wet or very dry may also result in poor weed control due to insufficient mixing. Refer to the General Instructions, especially “Cereal Seed Dressings” and “Weather Conditions at Planting” for further information on emergence.

SITUATION / CROP	WEEDS CONTROLLED	STATE
Cotton	See Weed Table A	Qld, NSW only
Soybeans, Peanuts, Navy Beans, Cow Peas, Mung Beans and Pigeon Peas		Qld, NSW, NT only
Sunflowers		Qld, NSW, Vic, SA only
Wheat and Barley	Annual Ryegrass (<i>Lolium rigidum</i>), Wireweed (<i>Polygonum aviculare</i>), Suppression of Wild Oats (<i>Avena</i> spp.)	All States
Field Peas		Southern NSW, Vic, Tas, SA, WA only
Chickpeas, Faba Beans, Lupins and Safflower		Qld, northern NSW, Tas* only

RATE / TREATED ha	CRITICAL COMMENTS
2.25 L/ha	Use a combine with trailing harrows and chain or offset or tandem discs or Lilliston cultivators, set to work to a depth of 2-5cm. Subsequent planting operations in irrigated cotton may remove the Genfarm Pendimethalin 440 EC Herbicide treated soil from hills leaving an untreated band over the row, requiring a further application of Genfarm Pendimethalin 440 EC Herbicide or other herbicides in that band. When the seedbed is rough, a second incorporation is necessary.
1.9 to 2.5 L/ha	Genfarm Pendimethalin 440 EC Herbicide may be applied by aerial or ground rig. In the Macquarie Valley area, only apply by air when ground is too wet for ground application.
900 ml/ha	Use weighted harrows at 8-13 km/hr followed by a cross working with a combine set to work at a depth of 2-5cm at 8-13 km/hr. Wheat and barley seed must be sown approximately 1cm beneath the chemically treated band or reduced emergence may occur. If Wild Oats are expected to be a problem use a Wild Oat herbicide.
1.5 L/ha	Use weighted harrows at 8-13 km/hour followed by a cross-working with a combine set to work at a depth of 2-5cm at 8-13 km/hr. Seed should be sown below the chemical band. Use cover harrows behind the combine. Use a Wild Oat herbicide, if Wild Oats are expected to be a problem.
1.9 to 2.5 L/ha	<p>*Lupins: In Tasmania, use only on the species <i>Lupinus angustifolius</i>. DO NOT use on Lalbus.</p> <p>Where full (double) incorporation is impractical and where the seedbed tilth is fine and free of large stones and trash, apply Genfarm Pendimethalin 440 EC Herbicide or in the case of lupins Genfarm Pendimethalin 440 EC Herbicide plus Simazine up to 24 hours prior to sowing. Incorporate with the combine set to sow to a depth of 3-5cm. Use trailing harrows. (Refer to “Incorporation” under the heading General Instructions). Application to heavier soil types that have poor structural characteristics or are poorly worked may result in insufficient and uneven incorporation and unsatisfactory weed control. Heavy soils that are cloddy or have a surface crust or surface seal should be avoided. Attempting to incorporate Genfarm Pendimethalin 440 EC Herbicide when soils are very wet or very dry may also result in insufficient mixing and therefore poor weed control. Refer to “Weather Conditions at Planting” under GENERAL INSTRUCTIONS, for further information on emergence.</p> <p>If Wild Oats are expected to be a problem use a Wild Oat herbicide.</p> <p>Use the higher rates on medium to heavy textured soils and the lower rate on light textured soils.</p>

SITUATION / CROP	WEEDS CONTROLLED	STATE
Wheat and Barley	Annual Ryegrass (<i>Lolium rigidum</i>), Wireweed (<i>Polygonum aviculare</i>), Suppression of Wild Oats (<i>Avena</i> spp.)	All States
Canola (Including crops with undersown Lucerne)	Suppression only of Silver Grass (<i>Vulpia</i> spp.)	Southern NSW, Vic, Tas, SA, WA only
Chickpeas, Faba Beans, Safflower, Lupins and Field Peas	Annual Ryegrass (<i>Lolium rigidum</i>), Wireweed (<i>Polygonum aviculare</i>), Suppression only of Silver Grass (<i>Vulpia</i> spp.) and Wild Oats (<i>Avena</i> spp.)	Southern NSW, Vic, SA, WA only
Lupins		WA only
		NSW, Vic, SA only

POST-PLANT PRE-EMERGENCE APPLICATION (OVERALL BAND SPRAYING): NON-MECHANICAL INCORPORATION

SITUATION / CROP	WEEDS CONTROLLED	STATE
Cotton	See Weed Table B	Qld, NSW only
Sunflowers		Qld, NSW, SA only
Maize		Qld, NSW, NT only

RATE / TREATED ha	CRITICAL COMMENTS
1.4 L/ha	<p>All crops (except Canola): Where full (double) incorporation is impractical and where the seedbed tilth is fine and free of large stones and trash, apply Genfarm Pendimethalin 440 EC Herbicide, or in the case of lupins Genfarm Pendimethalin 440 EC Herbicide plus Simazine, up to 24 hours prior to sowing. Incorporate with the combine set to sow to a depth of 3-5cm. Use trailing harrows. (Refer to “Incorporation” under the heading General Instructions). Wheat and barley seed must be sown at least 1 cm beneath the chemically treated band or reduced emergence may occur.</p> <p>If Wild Oats are expected to be a problem use a Wild Oat herbicide. Refer to Critical Comments on previous page plus:</p> <p>Canola: Apply Genfarm Pendimethalin 440 EC Herbicide up to 24 hours prior to sowing. Ensure Genfarm Pendimethalin 440 EC Herbicide is well incorporated. Sow Canola with a seeding depth of 1-3 cm, at the appropriate depth for conditions. Canola seed can be placed within the treated band.</p> <p>Use the higher rates on medium to heavy textured soils and the lower rates on light textured soils.</p> <p>Note: Development in waterlogged conditions after the application of Genfarm Pendimethalin 440 EC Herbicide may retard the emergence of Canola.</p> <p>If Wild Oats are expected to be a problem use a Wild Oat herbicide. Application to heavier soil types that have poor structural characteristics or are poorly worked may result in insufficient and uneven incorporation and unsatisfactory weed control. Heavy soils that are cloddy or have a surface crust or surface seal should be avoided. Attempting to incorporate Genfarm Pendimethalin 440 EC Herbicide when soils are very wet or very dry may also result in insufficient mixing and therefore poor weed control.</p> <p>Caution: Ensure thorough incorporation to maximise safety to undersown Lucerne. Use the lower rate when Canola is undersown with Lucerne.</p> <p>DO NOT apply to Canola post plant pre-emergence.</p>
1.4 to 2.25 L/ha	
1.5 to 2.25 L/ha	
1.5 to 2.25 L/ha	
1.5 L/ha plus 560 g/ha Simazine 900 WG or 1 L/ha Flowable Simazine	
1.5 L/ha plus 1.7 kg/ha Simazine 900 WG or 3 L/ha Flowable Simazine	

RATE / TREATED ha	CRITICAL COMMENTS
3.4 L/ha	Where incorporation prior to sowing is impractical and where the seedbed tilth is fine and free of large stones and trash, Genfarm Pendimethalin 440 EC Herbicide should be applied within 48 hours after sowing.
	Where incorporation prior to sowing is impractical and where the seedbed tilth is fine and free of large stones and trash, Genfarm Pendimethalin 440 EC Herbicide should be applied within 48 hours after sowing and before the emergence of the crop and weeds.
	Where Fat Hen or Blackberry Nightshade is expected to be a major problem, pre-plant incorporation is preferred. Refer to the relevant sections under the heading “Incorporation” in “General Instructions”.
3.4 L/ha	Use Genfarm Pendimethalin 440 EC Herbicide after planting and before emergence of crop or weeds, especially where grasses are expected to be the major problem.
	DO NOT incorporate. Avoid application in dry weather; this may cause pruning of aerial roots and lodging. Refer to the relevant sections under the heading “Incorporation” in “General Instructions”.

SITUATION / CROP	WEEDS CONTROLLED	STATE
Maize – <i>continued</i>	See Weed Table C	Qld, NSW, NT only
Broad Beans	Poppies, Wireweed (<i>Polygonum aviculare</i>)	Tas only
Carrots	See Weed Table D	Qld, NSW, Vic, SA, Tas, WA only
	Weed Table D plus Wild Radish (<i>Raphanus raphanistrum</i>) and Sowthistle (<i>Sonchus oleraceus</i>)	
Processing Peas	See Weed Table D	Qld, Vic only
		Tas only
French Beans		Qld, Tas only
Onions	Hogweed (Wireweed) (<i>Polygonum aviculare</i>)	Tas only

RATE / TREATED ha	CRITICAL COMMENTS
2.25 L/ha plus 1.1 kg/ha Atrazine 900 WG or 2 L/ha Flowable Atrazine	Genfarm Pendimethalin 440 EC Herbicide should be applied in a tank mix combination with atrazine after planting and before the emergence of crop and weeds, where Thornapple, Noogoora Burr, Bathurst Burr, Caltrop, Paddy Melon or Mintweed are expected to be a major problem. DO NOT incorporate this mixture. Follow directions on atrazine label carefully. This treatment should not lead to soil residue problems with atrazine, as the application rate is low. DO NOT use Genfarm Pendimethalin 440 EC Herbicide mixed with atrazine on heavy clay soils, as weed control may not be adequate. Refer to the relevant sections under the heading “Incorporation” in “General Instructions”.
3 L/ha	MINIMUM PLANTING DEPTH:
1.5 to 2.25 L/ha	Carrots: 15mm. Processing peas and French beans: 30mm.
1.5 L/ha plus linuron at 1.0kg ai/ha	Seed should be sown with press wheels or rollers behind the planter in order to ensure coverage of the seed and compaction of the seedbed. Apply Genfarm Pendimethalin 440 EC Herbicide to a fine seedbed free of ridges, clods and trash within 2 days after sowing.
1.5 to 3 L/ha	
3 L/ha	Use the higher rates on heavier textured soils or those with a higher organic matter content. Incorporate with 12 to 25mm of spray irrigation or rainfall applied within one day of application for optimum performance. DO NOT disturb the soil by rolling or harrowing after Genfarm Pendimethalin 440 EC Herbicide has been applied. DO NOT apply where heavy rainfall or irrigation is likely to lead to water logging. DO NOT use Genfarm Pendimethalin 440 EC Herbicide on soils with an organic matter content above 6% as inconsistent weed control will occur.
2.25 to 3 L/ha	French Beans: For use in the Lockyer and Fassifern Valleys only. DO NOT apply to French Beans sown during the autumn, winter or early spring or which are likely to be stressed by cold weather.
0.56 to 0.75 L/ha	Apply in conjunction with other herbicides as advised by a qualified crop adviser to ensure that an adequate level of weed control is achieved in adverse conditions. Pre-Emergence: DO NOT use on sandy soils. MINIMUM PLANTING DEPTH: 15mm. Apply from immediately after sowing until just prior to emergence. Apply Genfarm Pendimethalin 440 EC Herbicide to a fine, firm seedbed free of clods and trash. Use the higher rates on heavy textured soils or those with a high organic matter content. For optimum performance, with no more than 12 mm of spray irrigation within one day of application. DO NOT exceed 0.56 L/ha if soil conditions are wet and cold.
0.75 to 1.5 L/ha	Post-Emergence: Apply Genfarm Pendimethalin 440 EC Herbicide from the first true leaf until the 3 leaf stage. Repeat applications may be made after the 1 leaf up to the 3 leaf stage providing total usage of Genfarm Pendimethalin 440 EC Herbicide on crop does not exceed 2.25L/ha. Avoid applying Genfarm Pendimethalin 440 EC Herbicide to areas where water logging is likely to occur.

SITUATION / CROP	WEEDS CONTROLLED	STATE
Onions – <i>continued</i>	Hogweed (Wireweed) (<i>Polygonum aviculare</i>)	NSW, ACT, Vic, SA, WA only
		NSW, ACT, Vic, SA, WA only

RATE / TREATED ha	CRITICAL COMMENTS
	<ul style="list-style-type: none"> • Apply in conjunction with other herbicides as advised by a qualified crop advisor to ensure that an adequate level of weed control is achieved in adverse conditions. • Depth of sowing is critical. Onions must be sown to a depth of at least 15 mm with press wheels or rollers behind the planter to ensure coverage and compaction. • DO NOT disturb the soil by rolling or harrowing after application. • For optimum performance, incorporate with spray irrigation within a day of planting. However, DO NOT water heavily (no more than 12 mm irrigation) for at least 5 days after application. • AVOID application to areas where water logging is likely to occur. • DO NOT apply if heavy rain is likely within a week of application, as crop damage may result. • DO NOT exceed 0.75 L/ha if soil conditions are wet and cold. • DO NOT apply more than 3 times per season. • DO NOT apply more than 3 L/ha per season.
0.56 to 0.75 L/ha (See Critical Comments)	<p>Light/Sandy soils DO NOT apply pre-emergence.</p> <p>Application to moist soil is best Apply 0.56 to 0.75 L/ha post emergence at the 4-5 leaf stage, with 1-2 further applications of 0.56 to 0.75 L/ha as needed throughout the season. (Approx. 3-4 weeks apart).</p>
0.75 to 1.5 L/ha (See Critical Comments)	<p>Heavy/High organic matter soils Application to moist soil is best Apply 0.75 to 1.5 L/ha pre-emergence within 2 days of planting. Apply to a fine, firm seed bed free of ridges, clods and trash.</p> <p>For furrow irrigated crops, application to a dry seed bed, followed by irrigation within 5 days is optimal. An application of 0.75 to 1.5 L/ha post emergence may be made at the 2 leaf stage. (Application at this stage may cause yellowing of onions).</p>

POST-PLANT PRE-EMERGENCE TREATMENT (INCORPORATED OR NON-INCORPORATED)		
SITUATION / CROP	WEEDS CONTROLLED	STATE
Sugarcane	Summer Grass (<i>Digitaria ciliaris</i>), Awnless Barnyard Grass (<i>Echinochloa colona</i>), Crowsfoot Grass (<i>Eleusine indica</i>), Guinea Grass (<i>Panicum maximum</i> c.v. <i>Hamil</i> and c.v. <i>Guinea</i>), Green Summer Grass (<i>Brachiaria subquadriflora</i>)	Qld, NSW, WA only
	Blue Top (Blue Billygoat Weed) (<i>Ageratum houstonianum</i>)	
	Barnyard Grass (<i>Echinochloa</i> spp.), Blackberry Nightshade (<i>Solanum nigrum</i>), Blue Top (Blue Billygoat Weed) (<i>Ageratum houstonianum</i>), Common Sida (<i>Sida rhombifolia</i>), Giant/Black Pigweed (<i>Trianthema poertulacastrum</i>), Glossy Nightshade (<i>Solanum americanum</i>), Green Amaranth (<i>Amaranthus viridis</i>), <i>Ipomoea</i> spp., Liverseed Grass (<i>Urochloa panicoides</i>), Milkweed (Mexican Fire Paint) (<i>Euphorbia heterophylla</i>), Pigweed (<i>Portulaca oleracea</i>)	Qld, NSW, WA only
Combine sown rice	Barnyard Grass (<i>Echinochloa</i> spp.), Silvertop Grass (Brown Beetle Grass) (<i>Diplachne reptatrix</i>), Starfruit (<i>Damasonium minus</i>)	Qld, NSW, NT only

RATE / TREATED ha	CRITICAL COMMENTS
2.25 to 3.4 L/ha	Genfarm Pendimethalin 440 EC Herbicide should be applied as soon as possible after planting and before weed emergence. Consolidation (cutaway operation) of the planting furrow prior to application, to prevent soil movement into the base of the drill, is recommended. Expect poor grass control in soils that contain a high percentage of organic matter. Refer to the relevant sections under the heading “Incorporation” in “General Instructions”. Use the higher rate where longer weed control is required or where incorporation by irrigation or rainfall is likely to be delayed more than 5 days.
2.25 L/ha plus 1.7 kg/ha Atrazine 900 WG or 3 L/ha Flowable Atrazine or 1.7 kg/ha Diuron 900 WG	<p>NOTE INCORPORATION: Under dry conditions or where some cultivation is expected after application Genfarm Pendimethalin 440 EC Herbicide should be thoroughly incorporated by finger rakes to a shallow depth. In subsequent tillage operations, cultivation depth should not exceed the depth of incorporation.</p> <p>WATER INCORPORATION: Incorporate using 12 to 25mm of spray irrigation or when suitable rainfall is expected within 10 days of application.</p> <p>The area should not be disturbed by cultivation for the expected duration of weed control where this method of application is used. If excessive irrigation or rainfall occurs after the application of Genfarm Pendimethalin 440 EC Herbicide, resulting in the movement of soil from the top to the bottom of the planting drill, weed control will be reduced. For other broadleaved weeds higher rates of atrazine or diuron can be tank mixed with Genfarm Pendimethalin 440 EC Herbicide – see the relevant atrazine or diuron labels for details.</p> <p>TANK MIXTURES: Read and follow all label directions including restraints, spray drift restraints, mandatory no-spray zones, critical comments, withholding periods, regional use restrictions and safety directions for the tank mix products.</p>
2.25 L/ha plus 400 mL/ha Imazapic 240 g/L	Incorporate into the soil either by overhead irrigation, rainfall or mechanical incorporation, to a depth of 5 cm. Incorporate as soon as possible and within 3-5 days after application. Read all directions for use in Sugarcane on the Imazapic label.
2.25 L/ha plus propanil at 1.8 kg ai/ha	<p>Apply this tank mixture up to the 2 leaf stage of Barnyard Grass and 1 leaf stage of Silvertop, either by ground rig in 70-100 L/ha or by aircraft in 25-30 L/ha total spray volume.</p> <p>WATER MANAGEMENT: Flush immediately after sowing to enable a good even germination of rice and to ensure the soil surface is sealed. Apply a second flush or permanent flood after 2 days but not later than 5 days after the application. To assist in the control of Barnyard Grass, permanent water should be applied as soon as the rice can tolerate it and levels maintained for optimum growth of rice.</p> <p>RICE STAGE: After the first flushing irrigation, the stage of rice at application of this tank mixture is not critical. Timing should be based on the stage of weed growth and water management. However, DO NOT apply to rice in a weakened condition as a result of soil salts, moisture stress, over-watering, or any other cause. Some transient leaf burn of the rice may result after the application but no long term effects will result. Rice seed should be covered with at least 1cm of water after the first flushing irrigation before Genfarm Pendimethalin 440 EC Herbicide plus propanil is applied to germinating rice seed.</p>

POST-PLANT AND FLUSHING:		
SITUATION / CROP	WEEDS CONTROLLED	STATE
Combine sown rice and pasture sod sown rice	As above plus suppression of Annual Sedges (<i>Cyperus</i> spp.)	Qld, NSW, NT only

PRE-TRANSPLANT TREATMENT		
SITUATION / CROP	WEEDS CONTROLLED	STATE
Transplanted broccoli, cabbage and cauliflower	See Weed Table D	All States
Transplanted lettuce	See Weed Table D	Qld, NSW, Vic, SA, WA only

RATE / TREATED ha	CRITICAL COMMENTS
2.25 or 3.4 L/ha plus 800 mL/ha paraquat 250 g/L	<p>Use the higher rate when sowing into pasture sod sown rice for Annual Sedge control or when more than 10 days is expected between treatment and permanent water. Apply up to the 3 leaf stage of Barnyard Grass, Silvertop and Sedges, before rice emergence. Apply by ground rig in 70-100 L/ha of water total spray volume.</p> <p>WATER MANAGEMENT: Flush immediately after sowing to enable a good even germination of rice. The soil surface must be sealed by a flushing irrigation or rainfall before the application of Genfarm Pendimethalin 440 EC Herbicide plus paraquat. Apply a second flush or permanent flood after 1 day but not later than 5 days after the application of Genfarm Pendimethalin 440 EC Herbicide plus paraquat. To assist in the control of Barnyard Grass, permanent water should be applied as soon as the rice can tolerate it and levels maintained for optimum growth of rice.</p> <p>RICE STAGE: After the first flushing irrigation, Genfarm Pendimethalin 440 EC Herbicide plus paraquat must be applied before rice emergence. DO NOT apply to rice in a weakened condition from causes such as soil salts, moisture stress, over-watering or any other cause.</p>

RATE / TREATED ha	CRITICAL COMMENTS
1.5 to 2.25 L/ha	<p>Apply Genfarm Pendimethalin 440 EC Herbicide to a firm seedbed free of ridges, clods and trash between 2-7 days before transplanting. Use the higher rates on heavier textured soils or those with higher organic matter content. For optimum performance incorporate with 12 to 25mm of spray irrigation (or rainfall expected within one day) and ensure minimal mechanical disturbance to the seedbed at transplanting.</p> <p>DO NOT apply after transplanting, as damage will occur. DO NOT apply where water logging is likely to occur after transplanting or crop stunting will result. DO NOT use Genfarm Pendimethalin 440 EC Herbicide on soils with an organic matter content above 6% as inconsistent weed control will occur.</p> <p>Ramrod* may be tank mixed with Genfarm Pendimethalin 440 EC Herbicide prior to transplanting or applied separately after transplanting.</p>
1.5 to 3 L/ha	<p>DO NOT apply after transplanting, as damage will occur. DO NOT apply where water logging is likely to occur after transplanting or crop stunting will result. DO NOT use Genfarm Pendimethalin 440 EC Herbicide on soils with an organic matter content above 6% as inconsistent weed control will occur.</p> <p>Ramrod* may be tank mixed with Genfarm Pendimethalin 440 EC Herbicide prior to transplanting or applied separately after transplanting.</p>

PERENNIAL CROPS		
SITUATION / CROP	WEEDS CONTROLLED	STATE
Avocados, bananas, citrus, deciduous fruits, grapevines, lychees, macadamia nut, mangoes, nuts	Dwarf Amaranth (<i>Amaranthus macrocarpus</i>), Green Amaranth (<i>Amaranthus viridis</i>), Annual Ryegrass (<i>Lolium rigidum</i>), Asthma Plant (<i>Euphorbia hirta</i>), Barnyard Grass (<i>Echinochloa</i> spp.), Chickweed (<i>Stellaria media</i>), Crowsfoot Grass (<i>Eleusine indica</i>), Deadnettle (<i>Lamium amplexicaule</i>), Fat Hen (<i>Chenopodium album</i>), Pigeon Grass (<i>Setaria</i> spp.), Pigweed (<i>Portulaca oleracea</i>), Prickly Lettuce (<i>Lactuca serriola</i>), Sowthistle (<i>Sonchus oleraceus</i>), Summer Grass (<i>Digitaria anguinalis</i>), Winter Grass (<i>Poa annua</i>), Wireweed (<i>Polygonum aviculare</i>)	All States
	As above plus Prairie Grass (<i>Bromus unioloides</i>), Spotted Medic (<i>Medicago arabica</i>)	
Deciduous fruits, nuts, grapevines and citrus,	Asthma Plant (<i>Euphorbia hirta</i>), Summer Grass (<i>Digitaria anguinalis</i>), Barnyard Grass (<i>Echinochloa</i> spp.), Crowsfoot Grass (<i>Eleusine indica</i>), Pigeon Grass (<i>Setaria</i> spp.), Winter Grass (<i>Poa annua</i>), Annual Ryegrass (<i>Lolium rigidum</i>), Chickweed (<i>Stellaria media</i>), Sowthistle (<i>Sonchus oleraceus</i>), Pigweed (<i>Portulaca oleracea</i>), Dwarf Amaranth (<i>Amaranthus macrocarpus</i>), Green Amaranth (<i>Amaranthus viridis</i>), Fat Hen (<i>Chenopodium album</i>), Wireweed (<i>Polygonum aviculare</i>), Prickly Lettuce (<i>Lactuca serriola</i>), Deadnettle (<i>Lamium amplexicaule</i>)	
	As above plus Stinking Roger (<i>Tagetes minuta</i>), Cobbler's Pegs (<i>Bidens pilosa</i>), Curious Weed (<i>Schkuhria pinnata</i>)	
Deciduous fruits, nuts, grapevines and citrus	As above plus Caltrop (<i>Tribulus terrestris</i>)	All States
Established Lucerne seed crops only	Summer Grass (<i>Digitaria anguinalis</i>), Fat Hen (<i>Chenopodium album</i>)	NSW, SA only
	As above plus Pigeon Grass (<i>Setaria</i> spp.)	
Eucalypt forestry plantations	Amaranth (<i>Amaranthus</i> spp.), Annual Ryegrass (<i>Lolium rigidum</i>), Asthma Plant (<i>Euphorbia hirta</i>), Barnyard Grass (<i>Echinochloa</i> spp.), Chickweed (<i>Stellaria media</i>), Crowsfoot Grass (<i>Eleusine indica</i>), Deadnettle (<i>Lamium amplexicaule</i>), Fat Hen (<i>Chenopodium album</i>), Pigeon Grass (<i>Setaria</i> spp.), Pigweed (<i>Portulaca oleracea</i>), Prickly Lettuce (<i>Lactuca serriola</i>), Stagger Weed (<i>Stachys arvensis</i>), Sowthistle (<i>Sonchus oleraceus</i>), Summer Grass (<i>Digitaria anguinalis</i>), Winter Grass (<i>Poa annua</i>), Wireweed (<i>Polygonum aviculare</i>)	All States

RATE / TREATED ha	CRITICAL COMMENTS
6.75 to 9 L/ha	DO NOT apply Genfarm Pendimethalin 440 EC Herbicide to grapevines after bud swell. Precaution: Genfarm Pendimethalin 440 EC Herbicide or Genfarm Pendimethalin 440 EC Herbicide plus simazine is likely to affect the emergence and growth of green manure crops such as oats or beans between rows of trees or vines, if they are sown into the treated band within 12 months of application. Use the higher rate on medium to heavy textured soils and the lower rate on light textured soils. Use a directed spray avoiding spray contact with green bark, fruit and foliage. If applied to freshly transplanted trees or vines, soils should be compacted prior to application of Genfarm Pendimethalin 440 EC Herbicide to avoid contact with roots. Genfarm Pendimethalin 440 EC Herbicide must be incorporated by a minimum of 5mm of rainfall or spray irrigation as soon as possible but no later than 10 days after application or weed control may be reduced.
9 L/ha	For best results, soil surface should be free of weeds, surface litter and clods at the time of application. Genfarm Pendimethalin 440 EC Herbicide should be tank mixed with a knockdown herbicide at the recommended rate. Crop damage and/or reduced weed control may occur if Genfarm Pendimethalin 440 EC Herbicide is not incorporated successfully. If insufficient rainfall follows within 10 days of application the product must be incorporated over the whole surface area treated through the use of overhead sprinklers or by mechanical means, using appropriate equipment to avoid root damage. Drip irrigation is inadequate for this purpose.
6.75 L/ha plus 900 g/ha Simazine 900 WG or 1.6 L/ha Flowable Simazine	A minimum of 10-15 mm of rain or sprinkler irrigation is normally required to incorporate Genfarm Pendimethalin 440 EC Herbicide into the soil. For best results, soil surface should be free of weeds, surface litter and clods at the time of application. If small weeds are present at the time of application, Genfarm Pendimethalin 440 EC Herbicide should be tank mixed with a knockdown herbicide at the recommended rate.
9L/ha plus 1.1 kg/ha Simazine 900 WG or 2 L/ha Flowable Simazine	Use of simazine could lead to crop damage on vines younger than 3 years of age. Refer to the simazine label. DO NOT apply tank mixtures of Genfarm Pendimethalin 440 EC Herbicide plus simazine to alkaline soils.
2.25 L/ha	During the spring months prior to irrigation, the Lucerne stand should be grazed down hard to produce a low trash soil surface. Apply Genfarm Pendimethalin 440 EC Herbicide and incorporate treatment by 12-25mm of spray irrigation or when rainfall is expected within one day of application.
3.4 L/ha	
6.75 to 9 L/ha	PRE-PLANTING: Apply 7-2 days prior to transplanting. Use the higher rate on heavier textured soils or soils with high organic matter. Mechanical disturbance should be minimised during planting to optimise performance. POST-PLANTING: Apply immediately following planting. Before emergence of weeds. Repeat applications may be made as directed sprays to provide pre-emergence control of weeds. Emerged weeds should be controlled with carefully directed applications of a non-selective herbicide or over- the-top application of a selective herbicide. If applied to freshly transplanted trees, soil should be compacted prior to application to avoid contact with roots. Weed control may be reduced if rainfall or irrigation does not occur within 10 days of application. Flood irrigation may reduce control.

SITUATION / CROP	WEEDS CONTROLLED	STATE
Pyrethrum Pre-emergence crops	See Weed Table D	Tas only
Pyrethrum Post-emergence crops		
Pyrethrum – newly transplanted “splits”	Seedling docks (<i>Rumex</i> spp.) Suppression of the following weeds is also obtained: Capeweed (<i>Arctotheca calendula</i>), Storks Bill (<i>Erodium</i> spp.), Spear Thistle (<i>Cirsium vulgare</i>)	Tas only
Established pyrethrum (one year old or greater)		
Teatree (<i>Melaleuca alternifolia</i>)	Weed Table B plus Dwarf Amaranth (<i>Amaranthus macrocarpus</i>), Green Amaranth (<i>Amaranthus vindis</i>), Annual Ryegrass (<i>Lolium rigidum</i>), Barnyard Grass (<i>Echinochloa</i> spp.), Chickweed (<i>Stellaria media</i>), Crowsfoot Grass (<i>Eleusine indica</i>), Deadnettle (<i>Lamium amplexicaule</i>), Fat Hen (<i>Chenopodium album</i>), Pigeon Grass (<i>Setaria</i> spp.), Pigweed (<i>Portulaca oleracea</i>), Prickly Lettuce (<i>Lactuca serriola</i>), Sowthistle (<i>Sonchus oleraceus</i>), Summer Grass (<i>Digitaria anguinalis</i>), Winter Grass (<i>Poa annua</i>), Wireweed (<i>Polygonum aviculare</i>)	NSW, Qld, SA, NT only

ESTABLISHED TURF		
SITUATION / CROP	WEEDS CONTROLLED	STATE
TURF WARM SEASON SPECIES Bahia Grass, Buffalo Grass, Couch, Kikuyu, salt Water Couch and Zoysia Grass Turf	Summer Grass, (<i>Digitaria anguinalis</i>), Crowsfoot (Crab Grass), Winter Grass (<i>Poa annua</i>)	All States
Queensland Blue Couch, Hybrid Couch c.v. Tifdwarf		
TURF COOL SEASON SPECIES Kentucky Blue Grass		
Bent Grass, Perennial Ryegrass		

RATE / TREATED ha	CRITICAL COMMENTS
1.5 to 2.25 L/ha	Apply Genfarm Pendimethalin 440 EC Herbicide to seedbed within 10 days of sowing, prior to pyrethrum emergence. Apply at the lower rates on sandy soils. Incorporate with 12-25 mm of rainfall or irrigation within 1 day of application for optimum performance.
1.5 to 3.75 L/ha	Apply Genfarm Pendimethalin 440 EC Herbicide prior to weed germination at any stage or pyrethrum growth from 2 true leaf stage to pre-flowering, as advised by qualified crop advisor. Seek advice from qualified crop advisor before applying follow-up applications in any one year. Incorporate with 12-25 mm of rainfall or irrigation within 1 day of application for optimum performance.
3.75 L/ha	Apply immediately following transplanting before emergence of weeds. Repeat applications in any one year should only be made according to advice from crop advisers. Use lower rates on light soils. This product should be used in combination with other registered herbicides as recommended by crop advisers. Apply in autumn or spring prior to weed germination. Note existing weeds must be controlled with a knockdown herbicide, as Genfarm Pendimethalin 440 EC Herbicide will not control these weeds. Repeat applications in any one year should only be made according to advice from crop advisers. This product should be used in combination with other registered herbicides as recommended by crop advisers.
3.4 to 6.75 L/ha	DO NOT tank mix with simazine as crop damage may result. For best results the soil surface should be free of weeds, surface litter and clods at the time of application. PRE-TRANSPLANTING: Apply 7 to 2 days prior to transplanting seedlings. Use the higher rates on heavier textured soils or soils with higher organic matter. Mechanical disturbance should be minimised during planting to optimise performance. Avoid contact of treated soil with seedling roots. POST-PLANTING: Apply immediately following planting and pre-emergence of weeds. If applied to freshly transplanted trees, soils should be compacted prior to application of Genfarm Pendimethalin 440 EC Herbicide to avoid contact with roots. Weed control may be reduced if rainfall or irrigation does not occur within 10 days of application. Flood irrigation may reduce control.

RATE / TREATED ha	CRITICAL COMMENTS
3.4 L/ha or 34 mL/100m ²	Incorporate by 10-15mm of spray irrigation or when rainfall is expected within one day of application. Apply once every 10 weeks from early September to the end of February.
2.25 L/ha or 22.5 mL/100m ²	
3.4 L/ha or 34 mL/100m ²	The product should be incorporated by 10-15mm of spray irrigation or rainfall as soon as possible within one day of application.
2.25 L/ha or 22.5 mL/100m ²	Apply Genfarm Pendimethalin 440 EC Herbicide only once per year to control weeds in Perennial Ryegrass and Bent Grass.

IRRIGATION CHANNELS		
SITUATION / CROP	WEEDS CONTROLLED	STATE
Cotton Irrigation Banks, Channels and Drains	Barnyard Grass (<i>Echinochloa</i> spp.), Silvertop Grass (Brown Beetle Grass) (<i>Diplachne reptatrix</i>)	Qld, NSW only

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED

RATE / TREATED ha	CRITICAL COMMENTS
3.4 to 6.75 L/ha	Use the lower rate when short term control is required. Apply in late winter or early spring after regrading or clearing channels. The soil should be loose and free of large clods. If 25-50mm of rainfall has not fallen within 14 days of application the channel should be filled with water and allowed to stand for 1 day. The water in the channel should then be drained off and used to pre-irrigate cotton fields. DO NOT use water in channel to irrigate or pre-irrigate susceptible crops.
4.5 L/ha plus 9 kg/ha Diuron 900 WG or 16.2 L/ha Flowable Diuron	Use this mixture where long term control is required or a broader weed spectrum is present. See diuron label for details. Water in the channel should NOT be used to irrigate or pre-irrigate susceptible crops. TANK MIXTURES: Read and follow all label directions including restraints, spray drift restraints, mandatory no-spray zones, critical comments, withholding periods, regional use restrictions and safety directions for the tank mix products.

WEED TABLE A	
IN COTTON, COW PEAS, MUNG BEANS, NAVY BEANS, PEANUTS, PIGEON PEAS, SOYBEANS, SUNFLOWERS – with pre-plant incorporated treatment (double incorporation)	
The following weeds are controlled:	
Barnyard Grass	<i>Echinochloa</i> spp.
Button Grass	<i>Dactyloctenium radulans</i>
Common Heliotrope	<i>Heliotropium europeum</i>
Crested Goosefoot	<i>Chenopodium cristatum</i>
Crowsfoot Grass	<i>Eleusine indica</i>
Dwarf Amaranth	<i>Amaranthus macrocarpus</i>
Early Spring Grass	<i>Eriochloa</i> spp.
Fat Hen	<i>Chenopodium album</i>
Green Amaranth	<i>Amaranthus viridis</i>
Liverseed Grass	<i>Urochloa panicoides</i>
Mexican Clover	<i>Richardia brasiliensis</i>
Mossman River Grass	<i>Cenchrus echinatus</i>
Native Millet	<i>Panicum decompositum</i>
Pale Pigeon Grass	<i>Setaria glauca</i>
Paspalidium	<i>Paspalidium</i> spp.
Pepper Grass	<i>Panicum whitei</i>
Pigweed	<i>Portulaca oleracea</i>
Queensland Blue Grass	<i>Dichanthium sericeum</i>
Red Flinders Grass	<i>Iseilema vaginiflorum</i>
Scarlet Pimpernel	<i>Anagallis arvensis</i>
Small Burr Grass	<i>Tragus australianus</i>
Stink Grass	<i>Eragrostis cilianensis</i>
Weeping Love Grass	<i>Eragrostis parviflora</i>
Wireweed	<i>Polygonum aviculare</i>
Useful suppression of the following weeds is also obtained:	
Blackberry Nightshade	<i>Solanum nigrum</i>
Caltrop	<i>Tribulus terrestris</i>
Common Verbena	<i>Verbena officinalis</i>
Mintweed	<i>Salvia reflexa</i>
Peppergrass	<i>Lepidium</i> spp.
Stagger Weed	<i>Stachys arvensis</i>

WEED TABLE B	
IN COTTON, SUNFLOWERS, MAIZE and TEA TREE – with post-plant pre-emergence surface treatment (Genfarm Pendimethalin 440 EC Herbicide alone)	
The following weeds are controlled:	
Barnyard Grass	<i>Echinochloa</i> spp.
Button Grass	<i>Dactyloctenium radulans</i>
Common Heliotrope	<i>Heliotropium europeum</i>
Crested Goosefoot	<i>Chenopodium cristatum</i>
Crowsfoot Grass	<i>Eleusine indica</i>
Dwarf Amaranth	<i>Amaranthus macrocarpus</i>
Early Spring Grass	<i>Eriochloa</i> spp.
Green Amaranth	<i>Amaranthus viridis</i>
Liverseed Grass	<i>Urochloa panicoides</i>
Mexican Clover	<i>Richardia brasiliensis</i>
Mossman River Grass	<i>Cenchrus echinatus</i>
Native Millet	<i>Panicum decompositum</i>
Pale Pigeon Grass	<i>Setaria glauca</i>
Paspalidium	<i>Paspalidium</i> spp.
Pepper Grass	<i>Panicum whitei</i>
Pigweed	<i>Portulaca oleracea</i>
Queensland Blue Grass	<i>Dichanthium sericeum</i>
Red Flinders Grass	<i>Iseilema vaginiflorum</i>
Scarlet Pimpernel	<i>Anagallis arvensis</i>
Small Burr Grass	<i>Tragus australianus</i>
Stink Grass	<i>Eragrostis cilianensis</i>
Weeping Love Grass	<i>Eragrostis parviflora</i>
Wireweed	<i>Polygonum aviculare</i>
Useful suppression of the following weeds is also obtained:	
Caltrop	<i>Tribulus terrestris</i>
Common Sowthistle	<i>Sonchus oleraceus</i>
Common verbena	<i>Verbena officinalis</i>
Mintweed	<i>Salvia reflexa</i>
Peppergrass	<i>Lepidium</i> spp.
Stagger Weed	<i>Stachys arvensis</i>

WEED TABLE C	
IN MAIZE – with post-plant pre-emergence surface treatment of Genfarm Pendimethalin 440 EC Herbicide plus tank mix with atrazine.	
The following weeds are controlled in addition to those controlled by Genfarm Pendimethalin 440 EC Herbicide alone:	
Bathurst Burr	<i>Xanthium spinosum</i>
Caltrop	<i>Tribulus terrestris</i>
Mintweed	<i>Salvia reflexa</i>
Noogoora burr	<i>Xanthium pungens</i>
Paddy Melon	<i>Cucumis</i> spp.
Thornapple	<i>Datura</i> spp.
The following weeds are not controlled by GENFARM PENDIMETHALIN 440 EC HERBICIDE or the mixture with atrazine:	
Bindweed	<i>Convolvulus</i> spp.
Wandering Jew	<i>Commelina</i> spp.
Nut Grass	<i>Cyperus rotundus</i>

WEED TABLE D	
IN CARROTS, PROCESSING PEAS, FRENCH BEANS, TRANSPLANTED CABBAGE, CAULIFLOWER, BROCCOLI AND LETTUCE – with pre-emergence surface treatment after planting or before transplanting as directed and incorporated by spray irrigation.	
The following weeds are controlled:	
Awnless Barnyard Grass	<i>Echinochloa colona</i>
Bittercress	<i>Coronopus didymus</i>
Bladder Ketmia	<i>Hibiscus trionum</i>
Chickweed	<i>Stellaria media</i>
Deadnettle	<i>Lamium amplexicaule</i>
Fat Hen	<i>Chenopodium album</i>
Green Fat Hen	<i>Chenopodium murale</i>
Pigweed	<i>Portulaca oleracea</i>
Prickly Lettuce	<i>Lactuca serriola</i>
Shepherd's Purse	<i>Capsella bursa-pastoris</i>
Wireweed	<i>Polygonum aviculare</i>
Suppression only of the following weeds at the lower rates. Higher rates or tank mixtures are required for complete control (see Directions for Use table):	
Annual Nettles	<i>Urtica</i> spp.
Blackberry Nightshade	<i>Solanum nigrum</i>
Common Sowthistle	<i>Sonchus oleraceus</i>
Fumitory	<i>Fumaria</i> spp.
Indian Hedge Mustard	<i>Sisymbrium orientale</i>
Turnip Weed	<i>Rapistrum rugosum</i>
Wild Radish	<i>Raphanus raphanistrum</i>
Winter Grass	<i>Poa annua</i>

GENERAL INSTRUCTIONS

Genfarm Pendimethalin 440 EC Herbicide is a selective herbicide for the control of most annual grasses and certain broadleaved weeds. Genfarm Pendimethalin 440 EC Herbicide controls weeds by inhibiting seedling development; it will not control established weeds. For best results, seedbeds should be free of weeds, trash and clods at the time of application. Soils containing a high percentage of organic matter can result in poor control. It is recommended that application to soils containing a high percentage of organic matter (greater than 6%) be avoided.

Crop injury may result if cold, wet weather follows planting or if Genfarm Pendimethalin 440 EC Herbicide is applied to a poorly prepared seedbed. With the return of favourable growing conditions (warm weather), the crop will usually recover and resume normal growth.

Note: Genfarm Pendimethalin 440 EC Herbicide will stain (yellow) any objects it contacts. Keep all objects, etc. that are not to be stained well clear of the product and spray.

MIXING

Put half the required volume of water in spray tank and start agitation. When using Genfarm Pendimethalin 440 EC Herbicide alone, add Genfarm Pendimethalin 440 EC Herbicide to partially filled tank and then add the remainder of the water. Mix thoroughly. When using Genfarm Pendimethalin 440 EC Herbicide in tank mixes with products such as atrazine or diuron, mix these products as indicated on the relevant label and add the Genfarm Pendimethalin 440 EC Herbicide last, then complete filling of the tank. Maintain good agitation at all times until spraying is completed. If the spray mixture is allowed to settle, thorough agitation is essential to re-suspend the mixture before spraying is resumed. Note that when tank mixing Genfarm Pendimethalin 440 EC Herbicide with paraquat and glyphosate herbicides, the Genfarm Pendimethalin 440 EC Herbicide should be thoroughly mixed in the spray tank first, before adding a paraquat or glyphosate product.

APPLICATION METHOD

Ground Application:

Use conventional sprayers with either mechanical or by-pass agitation. Flat fan nozzles should be used. Spray equipment should be correctly calibrated to ensure proper application. If applying Genfarm Pendimethalin 440 EC Herbicide in tank mix combination with atrazine or diuron, nozzle screens must be no finer than 50 mesh. Apply the recommended quantity of Genfarm Pendimethalin 440 EC Herbicide in 50-200 litres of water per hectare. Where pre-emergence surface treatment is used, the higher rates of water are recommended.

Aerial Application:

Pre-plant soybeans, cotton, sunflowers, peanuts, navy beans, cow peas, mung beans, pigeon peas and post-plant, post flushed combine sown rice only.

Apply the recommended quantity of Genfarm Pendimethalin 440 EC Herbicide in 25 60 litres of water per hectare. It is essential that the equipment is correctly calibrated and an even spray pattern is obtained. It is essential that markers be used and an accurate swath width is maintained.

DO NOT apply if wind speed is greater than 16 km/hour. Always spray with a cross wind in accordance with recommended flying practice. DO NOT apply Genfarm Pendimethalin 440 EC Herbicide by aerial spraying in eucalypt forestry situations.

Incorporation:

- Genfarm Pendimethalin 440 EC Herbicide works best if thoroughly mixed with soil either mechanically or by irrigation or rainfall. The aim of incorporation is to produce an even band of herbicide to intercept germinating weed seeds.
- Post-plant pre-emergence surface applications of Genfarm Pendimethalin 440 EC Herbicide alone, or as a tank mixture, are most effective in controlling weeds when adequate rainfall occurs or irrigation is applied within 7-10 days after treatment to achieve incorporation. If no rain or irrigation is received within 7-10 days, light mechanical incorporation is required for optimum weed control, except when used on maize, carrots and rice.
- Maize is tolerant to Genfarm Pendimethalin 440 EC Herbicide either alone or in mixtures with atrazine if the seed germinates below the chemically treated band. Maize, carrots and rice are NOT tolerant to Genfarm Pendimethalin 440 EC Herbicide if it is in direct contact with the seed.

Cereal Seed Dressings

Some fungicidal seed dressings can shorten coleoptile length and this can affect emergence particularly if seed is sown too deep. When these cereal seed dressings are used in Genfarm Pendimethalin 440 EC Herbicide treated areas it is very important to ensure there is good seed depth control and to sow just below (1cm) the chemically treated band. Consult your local Department of Agriculture Agronomist for further advice.

Weather Conditions at Planting

Crop injury may result if cold weather follows planting. With the return of favourable growing conditions (warm weather), crops usually recover and resume normal growth. In cereals, prolonged cold wet conditions during germination and emergence can result in wheat or barley coleoptiles remaining in the treated band for an extended period. This can reduce emergence.

RESISTANT WEEDS WARNING

Genfarm Pendimethalin 440 EC Herbicide is a member of the dinitroaniline group of herbicides. Genfarm Pendimethalin 440 EC Herbicide has the inhibition of tubulin formation mode of action. For weed resistance management, Genfarm Pendimethalin 440 EC Herbicide is a Group 3 herbicide. Some naturally-occurring weed biotypes resistant to Genfarm Pendimethalin 440 EC Herbicide and other Group 3 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Genfarm Pendimethalin 440 EC Herbicide or other Group 3 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nutrien Ag Solutions Limited accepts no liability for any losses that may result from the failure of Genfarm Pendimethalin 440 EC Herbicide to control resistant weeds.

GROUP 3	HERBICIDE
----------------	------------------

COMPATIBILITY

Genfarm Pendimethalin 440 EC Herbicide is compatible with the following herbicides: atrazine, diuron, simazine, prometryn, fluometuron, fluometuron plus prometryn, ametryn, paraquat, propanil, linuron, glyphosate and propachlor.

If tank mixing, observe the directions under mixing.

FOLLOWING CROPS

In the event of crop failure the soil should be cultivated to a minimum depth of 15cm to ensure any residues are evenly dispersed throughout the soil.

If a maize crop fails due to weather conditions or some other reason, maize can be replanted the same year without adverse effects but seeding depth must be below the re-tilled area.

The following crops may be sown after a minimum interval of 2 months after application of Genfarm Pendimethalin 440 EC Herbicide: Carrots, parsnips, parsley, celery, potatoes, peas, French beans, transplanted Brassica's, transplanted lettuce, transplanted tomatoes, transplanted capsicums.

The following crops may be sown after a minimum interval of 5 months after application of Genfarm Pendimethalin 440 EC Herbicide: Turnips, lettuce, radish, Brussels sprouts, cabbage, cauliflower, broccoli, onions, leeks, Sweetcorn, pumpkins, squash, melons, cucumbers.

The following crops may be sown after a minimum interval of 12 months after application of Genfarm Pendimethalin 440 EC Herbicide: Redbeet (beetroot), spinach, silverbeet, poppy.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment, which could be expected to cause spray to drift onto nearby susceptible plants/adjacent crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and other aquatic life. DO NOT contaminate dams, rivers, drains or streams with chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store below 5°C. Extended storage below 5°C can result in the formation of crystals on the bottom of the container. If crystallisation does occur, store the container on its side at room temperature and rock occasionally until crystal re-dissolve. This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. DO NOT dispose of undiluted chemical on-site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection or similar container management program site. The cap should not be replaced, but may be taken separately. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

Refillable Containers: Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. DO NOT inhale vapour. When preparing spray and using the prepared spray. Wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length nitrile gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed, do NOT induce vomiting. If in eyes wash out immediately with water.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from the supplier.

CONDITIONS OF SALE: The use of this product is beyond the control of Nutrien Ag Solutions Limited. Any provisions or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Nutrien Ag Solutions Limited does not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: May be fatal if swallowed and enters airways. Very toxic to aquatic life with long lasting effects. Harmful in contact with skin. May cause an allergic skin reaction. Avoid breathing dust/fume/gas/mist/vapours/spray. Contaminated work clothing should not be allowed out of the workplace. Avoid release to the environment. Wear protective gloves / protective clothing. IF SWALLOWED: Immediately call a POISON CENTER/ doctor. Specific treatment (see on this label). Do NOT induce vomiting. IF ON SKIN: Wash with plenty of water. Call a POISON CENTER/doctor if you feel unwell. Take off contaminated clothing and wash it before reuse. If skin irritation or rash occurs: Get medical advice/attention. Collect spillage. Store locked up. Dispose of contents/container in accordance with local/regional/national regulations. Product contains: Pendimethalin (ISO), Liquid hydrocarbon



Nutrien Ag Solutions Limited
Level 5, Building A, 26 Talavera Road, Macquarie Park NSW 2113
Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92

APVMA Approval No: 91417/132175

10080041 0223